

## MICROPROPAGATION AND FLOWERING OF *GLORIOSA ROTHSCILDIANA* O'BRIEN WITH EFFICIENT THERMAL TREATMENT OF MICROTUBERS

Masao Yamamoto<sup>1,2</sup> and Masahiro Mii<sup>1\*</sup>

<sup>1</sup>Graduate School of Horticulture, Chiba University, 648 Matsudo, Matsudo, Chiba 271-8510, Japan,

\*Fax: + 81 47 308 8852, \*E-mail: miim@faculty.chiba-u.jp

<sup>2</sup>The Hiroshima Botanical Garden, 495, Kurashige 3-chome, Saekiku, Hiroshima, Hiroshima 731-5156, Japan

### Abstract

Procedures for the micropropagation and further *in vivo* cultivation of *Gloriosa rothschildiana* microtubers were established. Meristem and young shoot basal explants of *G. rothschildiana* were cultured on MS medium supplemented with 4.0 mg l<sup>-1</sup> BA and 0.1 mg l<sup>-1</sup> NAA, which resulted in the production of 5.0 and 47.2 mean numbers of shoots per explant, respectively, after 150 days of culture. Twice to 3 times of shoot multiplication was successfully achieved by subculturing the shoots onto the same medium. Increasing subculture duration on the same medium stimulated microtuber formation at the bottom of most shoots. Approximately 85% of the microtubers produced with a weight over 1 g sprouted successfully in *in vivo* conditions by the thermal treatment of 8°C for 2 weeks, followed by 30°C for 13 weeks. Sprouted tubers planted in soil grew normally and 80% of the tubers with weight over 5 g collected after 2 cycles of cultivation successfully produced flowers. No noticeable change was observed in the morphology including flower characters of the micropropagated plants.

**Key words:** acclimatization, *Gloriosa rothschildiana*, micropropagation, microtuber, ornamental flower, thermal treatment

### REFERENCES

- AKITA M., TAKAYAMA S. (1993). Resting period and the field performance of potato tubers propagated in a jar fermentor. *Plant Tissue Culture Letters*, 10 (3): 255-259.
- AKITA M., TAKAYAMA S. (1994). Stimulation of potato (*Solanum tuberosum* L.) tuberization by semicontinuous liquid medium surface level control. *Plant Cell Reports*, 13: 184-187.
- CUSTERS J. B. M., BERGERVOET J. H. W. (1994). Micropropagation of *Gloriosa*: Towards a practical protocol. *Scientia Horticulturae*, 57: 323-334.
- DE HERTOOGH A., LE NARD M. (1993). *The Physiology of Flower Bulbs*. Elsevier, Amsterdam: 755-756.
- ERHARDT W., GÖTZ E., BÖDEKER N., SEYBOLD S. (2002). *Zander: Handwörterbuch der Pflanzennamen*. Eugen Ulmer GmbH & Co., Stuttgart, 450 pp.
- GHOSH S., GHOSH B., JHA S. (2007). *In vitro* tuberisation of *Gloriosa superba* L. on basal medium. *Scientia Horticulturae*, 114: 220-223.
- KAWASHIMA T. (1983). The relationships between flowering and fresh weight of tubers of *Gloriosa rothschildiana*. *Annual Report of Hiroshima City Horticulture Institute*: 117-119 (in Japanese).
- KOZAK D. (2002). Studies on micropropagation of gloriosa (*Gloriosa rothschildiana* O'Brien). Dissertation of Agricultural University in Lublin, 261, 72 pp. (in Slovene).
- MURASHIGE T., SKOOG F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- SAMARAJEWA P. K., DASSANAYAKE M. D., JAYAWARDENA S. D. G. (1993). Clonal propagation of *Gloriosa superba* L. *Indian Journal of Experimental Biology*, 31: 719-720.
- SIVAKUMAR G., KRISHNAMURTHY K. V. (2004). *In vitro* organogenetic responses of *Gloriosa superba*. *Russian Journal of Plant Physiology*, 51: 790-798.
- YAMAMOTO K. (1990). Studies of thermal treatment for sprouting of dormant tubers of *Gloriosa rothschildiana*. *Annual Report of Hiroshima City Horticulture Institute*: 25-27 (in Japanese).