

**IN VITRO MORPHOGENESIS AND PLANT REGENERATION FROM SEED AND
LEAF EXPLANTS OF FOUR *AGROPYRON* SPECIES**

**Masoume Amirkhani^{1*}, Maurizio Lambardi², Mansour Mesdaghi¹,
Kambiz Mashayekhi³, and Elif Aylin Ozudogru⁴**

¹Range Management Department, Gorgan University of Agricultural Sciences and Natural Resources,
Behesht Avenue, Gorgan, Iran, *Fax: + 981 71 4470112, *E-mail: m.amirkhani1980@yahoo.com

²IVALSA, Trees and Timber Institute, National Research Council (CNR), via Madonna del Piano 10,
50019 Sesto Fiorentino, Firenze, Italy

³Horticulture Department, Gorgan University of Agricultural Sciences and Natural Resources,
Beheshti Avenue, Gorgan, Iran

⁴Gebze Institute of Technology, Faculty of Science, Biology Department, 101 Istanbul Caddesi,
41400 Gebze-Kocaeli, Turkey

REFERENCES

- ANGRA D. C., BARBOSA M. M., PRESTES A. M., FERNANDES M. (1999). Embryo culture of backcrosses between hybrids of *Triticum aestivum* Thell. and *Agropyron elongatum* Host. & Beauv. Pesquisa Agropecuaria Brasileira, 34 (2): 209-215.
- BARAMPURAM S., CHUNG B. Y., LEE S. S., CHULLAN B., LEE E. M., CHO J. Y. (2009). Development of an embryogenic callus induction method for centipede grass (*Eremochloa ophiuroides* Munro) and subsequent plant regeneration. In Vitro Cellular and Developmental Biology-Plant, 45:155-161.
- BHASKARAN S., SMITH R. H. (1990). Regeneration in cereal tissue culture: a review. Crop Science, 30: 1328-1336.
- BOR N. L. (1970). Gramineae, Tribus VII. Triticeae Dumort. In: Rechinger K. H. (Ed.). Flora Iranica, vol. 70, Verlagsanstalt, Graz, Austria: 147-244.
- CAN E., CELIKTAS N., HATIPOGLU R. (2008). Effect of auxin type and concentrations in different media on the callus induction and shoot formation of crested wheatgrass (*Agropyron cristatum* (L.) Gaertn). Biotechnology & Biotechnological Equipment, 22: 782-786.
- CHAUDHURY A., QU R. (2000). Somatic embryogenesis and plant regeneration of turf-type Bermuda grass: effect of 6-benzyladenine in callus induction medium. Plant Cell, Tissue and Organ Culture, 60:113-120.
- CHEKUROV V. M., RAZMAKHIN E. P. (1999). Effect of inbreeding and growth regulators on the *in vitro* androgenesis of wheatgrass, *Agropyron glaucum*. Plant Breeding, 118 (6): 571-573.
- CHEN Q., JAHIER J., CAUDERON Y. (1992). Production of embryo-callus-regenerated hybrids between *Triticum aestivum* and *Agropyron cristatum* possessing one B chromosome. Agronomie, 12: 551-555.
- DEWEY D. R. (1978). Intermediate wheat grasses of Iran. Crop Science, 18: 43-48.
- DEWEY D. R., ASAY K. H. (1975). The crested wheatgrass of Iran. Crop Science, 15: 844-849.
- DILMAN A. C. (1946). The beginnings of crested wheatgrass in North America. Journal of the American Society of Agronomy, 38(3): 237-250.
- GAMBORG O. L. (1991). Media preparation. In: Plant Tissue Culture Manual. Kluwer Academic Publishers, The Netherlands: 1-24.
- GRANDO F. M., FRANKLIN C. I., SHATTERS R. G. (2002). Optimizing embryogenic callus production and plant regeneration from 'Tifton 9' Bahia grass seed explants for genetic manipulation. Plant Cell, Tissue and Organ Culture, 71: 213-222.
- GUPTA S., KHANNA V. K., SINGH R., GARG C. K. (2006). Strategies for overcoming genotypic limitations of *in vitro* regeneration and determination of genetic components of variability of plant regeneration traits in sorghum. Plant Cell, Tissue and Organ Culture, 86: 379- 388.
- HUO X.-W., WEI J.-H., XU C.-B., MI F.-G., YUN J.-F. (2006). Plant regeneration from immature inflorescence culture and genetic transformation of wheatgrass (*Agropyron cristatum* × *A. desertorum* cv. Hycrest-Mengnong). Agricultural Science in China, 5 (9): 648-654.
- JIMENEZ V. M. (2001). Regulation of *in vitro* somatic embryogenesis with emphasis on the role of endogenous hormones. Revista Brasileira de Fisiologia Vegetale, 13:196-223.
- LATHA R., AJITH A., SRINIVASA R. C., EGANATHAN P., BALAKRISHNA P. (1998). *In vitro* micropropagation of salt tolerant wild rice relative, *Porteresia coarctata* Tateoka. Journal of Plant Growth Regulation, 17: 231-235.
- LI R., BRUNEAU A. H., QU R. (2006). Improved plant regeneration and *in vitro* somatic embryogenesis of St Augustine grass (*Stenotaphrum secundatum* (Walt.) Kunze). Plant Breeding, 125 :52-56.
- LO P. F., CHEN C. H., ROSS J. G. (1980). Vegetative propagation of temperate forage grasses through callus culture. Crop Science, 20: 363-367.
- MUJEEP-KAZI A., SUH D. Y., TER-KUILE N., FAROOQ S. (1989). Production and cytogenetics of *Triticum aestivum* L. hybrids with some rhizomatous *Agropyron* species. Theoretical and Applied Genetics, 77: 162-168.
- MURASHIGE T., SKOOG F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. Physiologia Plantarum, 15: 473-497.
- NEUMANN K. H. (1966). Wurzelbildung und Nukleinsäuregehalt bei Phloem-Gewebekulturen der Karottenwurzel auf synthetischen

- Nährmedien verschiedener Hormonkombinationen. Les Congres et Colloques de l'Universite de Liege, 38: 96-102.
- SAXENA S. (1990). *In vitro* propagation of the bamboo (*Bambusa tulda* Roxb.) through shoot proliferation. Plant Cell Reports, 9: 431-434.
- SHARMA H. C., GILL B. S., UYEMOTO J. K. (1984). High levels of resistance in *Agropyron* species to barley yellow dwarf and wheat streak mosaic virus. Phytopathologische Zeitschrift, 110: 143-147.
- SHOYAMA Y., ZHU X. X., NAKAI R., SHIRAISHI S., KOHDA H. (1997). Micropropagation of *Panax notoginseng* by somatic embryogenesis and RAPD analysis of regenerated plantlets. Plant Cell Reports, 16: 450-453.
- SPANGENBERG G., WANG Z. Y., POTRYKUS I. (1998). Biotechnology in Forage and Turf Grasses Improvement. Springer-Verlag, Berlin, Heidelberg, New York, 200 pp.
- WANG Z. Y., BELL J., HOPKIN A. (2003). Establishment of a plant regeneration system for wheatgrasses (*Thinopyrom*, *Agropyron* and *Pascopyrum*). Plant Cell, Tissue and Organ Culture, 73: 265-273.
- YAO K., CHEN C. H., McMULLEN C. R. (1991). Regeneration of albino shoots in cultured anthers of intermediate wheatgrass (*Agropyron intermedium* (Host) Beauv.). Plant Cell, Tissue and Organ Culture, 26: 189-193.