Propagation of Ornamental Plants
Vol. 13, № 1, 2013: 3-11

# MICRONUTRIENT OPTIMIZATION IN THE BASAL MEDIUM IMPROVES *IN VITRO* PLANT REGENERATION IN CARNATION (*DIANTHUS CARYOPHYLLUS* L.) AND CHINESE PINK (*DIANTHUS CHINENSIS* L.)

**Suman Kumawat, Sumita Kachhwaha, and Shanker Lal Kothari***

Department of Botany, University of Rajasthan, Jaipur, 302004, India,
*Fax: + 141-2703439, *E-mail: slkothari28@gmail.com

## REFERENCES

Amarasinghe A. A. Y. (2009). Effect of copper sulphate and cobalt chloride on *in vitro* performance of traditional Indica rice (*Oryza sativa* L.) varieties in Srilanka. Journal of Agricultural Science, 4: 132-141.

Bregitzer P., Campbell R. D., Dahleen L. S., Lemaux P. G., Cho M-J. (2000). Development of transformation systems for elite barley cultivars. Barley Genetics Newsletter, 30: 1-4.

Chauhan M., Kothari S. L. (2004). Optimization of nutrient levels in the medium increase the efficiency of callus induction and plant regeneration in recalcitrant Indian barley (*Hordeum vulgare* L.) *in vitro*. In Vitro Cellular & Developmental Biology-Plant, 40: 520-527.

Casanova E., Moysset L., Trillas M. I. (2008). Effects of agar concentration and vessel closure on the organogenesis and hyperhydricity of adventitious carnation shoots. Biologia Plantarum, 52: 1-8.

Casas J. L., Olmos E., Piqueras A. (2010). *In vitro* propagation of carnation (*Dianthus caryophyllus* L.). Methods in Molecular Biology, 589: 109-116.

Dahleen L. S., Bregitzer P. (2002). An improved media system for high regeneration rates from barley immature embryo-derived callus cultures of commercial cultivars. Crop Sciences, 42: 934-938.

Dangi B., Kachhwaha S., Kothari S. L. (2012). Regeneration and Agrobacterium-mediated genetic transformation of *Terminalia bellerica* Roxb.: a multipurpose tree species. In Vitro Cellular & Developmental Biology-Plant, 48: 304-312.

Debergh P., Aitken-Christie J., Cohen D., Grout B., Von Arnold S., Zimmetman R., Ziv M. (1992). Reconsideration of the term 'vitrification' as used in micropropagation. Plant Cell, Tissue and Organ Culture, 30:135-140.

Doyle J. J., Doyle J. L. (1990). Isolation of plant DNA from fresh tissue. Focus, 12: 13-15.

Earle E. D., Langhans R. W. (1975). Carnation propagation from shoot tips cultured in liquid medium. Horticultural Science, 10: 608-610.

Fatima N., Ahmad N., Anis M. (2011). Enhanced *in vitro* regeneration and change in photosynthetic pigments, biomass and proline content in *Withania somnifera* L. (Dunal) induced by copper and zinc ions. Plant Physiology and Biochemistry, 49: 1465-1471.

Frey L., Janick J. (1991). Organogenesis in carnation. Journal of the American Society for Horticultural Science, 116: 1108-1112.

Gatierrez-Miceli F. A., Arias L., Juarez-Rodriguez N., Abud-Archila M., Amaro-Reyes A., Dendooven L. (2010). Optimization of growth regulators and silver nitrate for micropropagation of *Dianthus caryophyllus* L. with the aid of a response surface experimental design. In Vitro Cellular & Developmental Biology-Plant, 46: 57-63.

Ghnaya A. B., Hourmant A., Cerantola S., Kervarec N., Cabon J. Y., Branchard M., Charles G. (2010). Influence of zinc on soluble carbohydrate and free amino acid levels in rapeseed plants regenerated *in vitro* in the presence of zinc. Plant Cell, Tissue and Organ Culture, 102: 191-197.

Gomez K. A., Gomez A. A. (1984). Statistical procedure for agriculture research. Second edition, John Wiley, New York, 680 pp.

Jain A., Kantia A., Kothari S. L. (2001). *De novo* differentiation of shoot buds from leaf callus of *Dianthus caryophyllus* L. and control of hyperhydricity. Scientia Horticulture, 87: 319-326.

Jain P., Kachhwaha S., Kothari S. L. (2009). Improved micropropagation protocol and enhancement in biomass and chlorophyll content in *Stevia rebaudiana* (Bert.) Bertoni by using high copper levels in the culture medium. Scientia Horticulture, 119: 315-319.

Jain R., Sinha A., Jain D., Kachhwaha S., Kothari S. L. (2011). Adventitious shoot regeneration and *in vitro* biosynthesis of steroidal lactones in *Withania coagulans* (stock) Dunal. Plant Cell, Tissue and Organ Culture, 105: 135-140.

Jethwani V., Kothari S. L. (1996). Phenylacetic acid induced organogenesis in cultured leaf segments of *Dianthus chinensis*. Plant Cell Report, 15: 869-872.

Joshi A., Kothari S. L. (2007). High cooper levels in the medium improves shoot bud differentiation and elongation from the cultured cotyledons of *Capsicum annuum* L. Plant Cell, Tissue and Organ Culture, 88: 127-133.

Kantia A., Kothari S. L. (2002). High efficiency adventitious shoot bud formation and plant regeneration from leaf explants of *Dianthus chinensis* L. Scientia Horticulture, 96: 205-212.

Kanwar J. K., Kumar S. (2009). Influence of growth regulators and explants on shoot regeneration in carnation. Horticultural Science, 36: 140-146.

Karami O., Karami-Kordestani G. (2007). Proliferation, shoot organogenesis and somatic embryogenesis in embryogenic callus of carnation. Journal of Fruit and Ornamental Plant Research, 15: 167-175.

Kevers C., Frank T., Strasser R. J., Dommes J., Gasper T. (2004). Hyperhydricity of micropropagated shoots: a typically stress-induced change of physiological state. Plant Cell, Tissue and Organ Culture, 77: 181-191.

Khurana-Kaul V., Kachhwaha S., Kothari S. L. (2010). Direct shoot regeneration from leaf explant of *Jatropha curcas* in response to thidiazuron and high copper content in the medium. Biologia Plantarum, 54: 369-372.

Kothari S. L., Agrawal K., Kumar S. (2004). Inorganic nutrient manipulation for highly improved *in vitro* plant regeneration in finger millet *Eleusine coracana* (L.) Gaertn. In Vitro Cellular & Developmental Biology-Plant, 40: 515-519.

Kothari-Chajer A., Sharma M., Kachhwaha S., Kothari S. L. (2008). Micronutrient optimization results in to highly improved *in vitro* plant regeneration in Kodo (*Paspalum scrobiculatum* L.) and Finger (*Eleusine coracana* L. Gaertn) millets. Plant Cell, Tissue and Organ Culture, 94: 105-112.

Kumar S., Narula A., Sharma M. P., Srivastava P. S. (2003). Effect of copper and zinc on growth, secondary metabolite content and micropropagation of *Tinospora cordiofolia*: A medicinal plant. Phytomorphology, 53: 79-91.

Kurniawan B. (2009). The occurrence of hyperhydricity on several carnation cultivars during low temperature storage. Biodiversitas, 10: 104-107.

Maksymiec W. (1997). Effect of cooper on cellular processes in higher plants. Photosynthetica, 34: 321-342.

Martins M., Sarmento D., Oliveira M. M. (2004). Genetic stability of micropropagated almond plantlets as assessed by RAPD and ISSR markers. Plant Cell Reports, 23: 492-496.

Masoni A. (1996). Special properties of leaves deficient in iron, sulfar, magnesium and magnese. Agronomy Journal, 88: 937-943.

Messeguer J., Arconada M. C., Mele E. (1993). Adventitious shoot regeneration in carnation (*Dianthus caryophyllus* L.). Scientia Horticulturae, 54: 153-163.

Miller R. M., Kaul V., Hutchinson J. F., Richards D. (1991). Adventitious shoot regeneration in carnation (*Dianthus caryophyllus*) from axillary bud explants. Annals of Botany, 67: 35-42.

Mouhtaridou G., Sotiropoulos T., Dimassi K., Therios I. (2004). Effect of boron on growth, and chlorophyll and mineral contents of shoots of the apple rootstock MM106, cultured *in vitro*. Biologia Plantarum, 48: 617-619.

Murashige T., Skoog F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiologia Plantarum, 15: 473-497.

Niedz R. P., Evens T. J. (2007). Regulating plant tissue growth by mineral nutrition. In Vitro Cellular & Developmental Biology-Plant, 43: 370-381.

Nirwan R. S., Kothari S. L. (2003). High copper levels improve callus induction and plant regeneration in *Sorghum bicolor* (L.) Moench. In Vitro Cellular & Developmental Biology-Plant, 39: 161-164.

Pareek A., Kantia A., Kothari S. L. (2004). *In vitro* cloning of ornamental species of *Dianthus*. Indian Journal of Biotechnology, 3: 263-266.

Poddar K., Vishnoi R. K., Kothari S. L. (1997). Plant regeneration from embryogenic callus of finger millets *Eleusine coracana* (L.) Gaertn. on higher concentration of $NH_4NO_4$ as a replacement of NAA in the medium. Plant Sciences, 129:101-106.

Prasad H. M. K., Mythili J. B., Tejaswini Lalitha Anand Rashmi H. J., Suneetha C. (2009). Optimization of regeneration protocol and *Agrobacterium* mediated transformation in carnation (*Dianthus caryophyllus* L.). Journal of Horticultural Sciences, 4: 120-127.

Purnhauser L., Gyulai G. (1993). Effect of copper on shoot and root regeneration in wheat, triticale, rape and tobacco tissue cultures. Plant Cell, Tissue and Organ Culture, 35: 131-139.

Ramage C. M., Williams R. R. (2002). Mineral nutrition and plant morphogenesis. In vitro Cellular & Developmental Biology-Plant, 38: 116-124.

Rani V., Raina S. N. (2000). Genetic fidelity of organized meristem-derived micropropagated plants: a critical reappraisal. In Vitro Cellular & Developmental Biology-Plant, 36: 319-330.

Rojas-Martínez L., Visser R G. F., De Klerk G-J. (2010). The hyperhydricity syndrome: waterlogging of plant tissues as a major cause. Propagation of Ornamental Plants, 10: 169-175.

Santandrea G., Schiff S., Bennici A. (1997). Manganese toxicity to different growth processes *in vitro* in *Nicotiana*. Plant Cell, Tissue and Organ Culture, 50:125-129.

Sharma A., Kumar V., Giridhar P., Ravishankar G. A. (2008). Induction of *in vitro* flowering in *Capsicum frutescens* under the influence of silver nitrate and cobalt chloride and pollen transformation. Electronic Journal of Biotechnology, 11: 84-89.

Sotiropoulos T. E., Dimassi K. N., Tsirakoglou V. (2006). Effect of boron and methionine on growth and ion content in kiwifruit shoots cultured *in vitro*. Biologia Plantarum, 50: 300-320.

Tahiliani S., Kothari S. L. (2004). Increased copper content of the medium improves plant regeneration from immature embryo derived callus of wheat (*Triticum aestivum*). Journal of Plant Biochemistry and Biotechnology, 13: 85-88.

Thakur M., Sharma D. R., Sharma S. K. (2002). *In vitro* regeneration of carnation (*Dianthus caryophyllus* L.) plants resistant to culture filtrate of *Fusarium oxysporum* f.sp. *dianthi*. Plant Cell Report, 20: 825-828.

van Altvorst A. C., Yancheva S., Dons H. (1995). Cells within the nodal region of carnation shoots exhibits a high potential for adventitious shoot formation. Plant Cell, Tissue and Organ Culture, 40: 151-157.

Vera-Estrella R., Maria C., Bronwyn J. (2009). Zinc tolerance and accumulation in stable cell suspension cultures and *in vitro* regenerated plants of the emerging model plants *Arabidopsis halleri* (Brassicaceae). Planta, 229: 977-986.

Wang H., Liu R. L., Jin J. Y. (2009). Effect of zinc and soil moisture on photosynthetic rate and chlorophyll fluorescence parameters of maize. Biologia Plantarum, 53: 191-194.

Watad A. A., Ahroni A., Shejtman H., Nissim A., Vainstein A. (1996). Adventitious shoot formation from carnation stem segments: a comparision of different culture procedures. Scientia Horticulture, 65: 313-320.

Yadav M. K., Gaur A. K., Garg G. K. (2003). Development of suitable protocol to overcome hyperhydricity in carnation during micropropagation. Plant Cell, Tissue and Organ Culture, 72: 153-156.