Propagation of Ornamental Plants
Vol. 16, № 4, 2016: 130-136

ADVANTAGEOUS EFFECTS OF MILD ABIOTIC STRESSES IN LILY CULTURED *IN VITRO*

Naser Askari[1,2], Richard G. F. Visser[1], and Geert-Jan de Klerk[1*]

[1]Wageningen UR Plant Breeding, P. O. Box 16, 6700 AA Wageningen, The Netherlands,
*Fax: + 31 317 418094, *E-mail: geertjan.deklerk@wur.nl
[2]Department of Plant Sciences, University of Jiroft, P. O. Box 364, Jiroft, Iran

REFRENCES

Aguettaz P., Paffen A., Delvallée I., Van Der Linde P., De Klerk G. J. (1990). The development of dormancy in bulblets of *Lilium speciosum* generated *in vitro*. Plant Cell, Tissue and Organ Culture, 22: 167-172.

Askari N. (2016). Aspects of bulblet growth of lily *in vitro*. PhD thesis Wageningen University, Wageningen, The Netherlands, 130 pp.

Chang P. T., Randle W. M. (2004). Sodium chloride in nutrient solutions can affect onion growth and flavor development. HortScience, 39: 1416-1420.

De Klerk G. J. M., Gerrits M. M. (1996). Development of dormancy in tissue-cultured lily bulblets and apple shoots. *In*: Lang G. A. (Ed.). Plant Dormancy - Physiology, Biochemistry and Molecular Biology, CAB International: 115-131.

De Klerk G. J., Pumisutapon P. (2008). Protection of *in-vitro* grown Arabidopsis seedlings against abiotic stresses. Plant Cell, Tissue and Organ Culture, 95: 149-154.

Delvallée I., Paffen A., De Klerk G. J. (1990). The development of dormancy in bulblets of *Lilium speciosum* generated *in vitro*. II. The effect of temperature. Physiologia Plantarum, 80: 431-436.

Dolferus R., Klok E. J., Delessert C., Wilson S., Ismond K. P., Good A. G., Peacock W. J., Dennis E. S. (2003). Enhancing the Anaerobic Response. Annals of Botany, 91: 111-117.

Ewing E. E. (1981). Heat stress and the tuberization stimulus. American Potato Journal, 58: 31-49.

Gong B., Yi J., Wu J., Sui J., Khan M., Wu Z., Zhong X., Seng S., He J., Yi M. (2014). LlHSFA1, a novel heat stress transcription factor in lily (*Lilium longiflorum*), can interact with LlHSFA2 and enhance the thermotolerance of transgenic *Arabidopsis thaliana*. Plant Cell Reports, 33: 1519-1533.

Hilker M., Schwachtje J., Baier M., Balazadeh S., Bäurle I., Geiselhardt S., Hincha D. K., Kunze R., Mueller-Roeber B., Rillig M. C., Rolff J., Romeis T., Schmülling T., Steppuhn A., Van Dongen J., Whitcomb S. J., Wurst S., Zuther E., Kopka J. (2016). Priming and memory of stress responses in organisms lacking a nervous system. Biological Reviews, 91: 1118-1133.

Jackson M. B. (1985). Ethylene and responses of plants to soil waterlogging and submergence. Annual Review of Plant Physiology and Plant Molecular Biology, 36: 145-174.

Langens-Gerrits M. M, De Klerk G. J., Croes A. (2000). How to produce lily bulblets *in vitro* that perform optimally after planting. Acta Horticulture, 530: 289-296.

Langens-Gerrits M., Kuijpers A. M., De Klerk G. J., Croes A. (2003). Contribution of explant carbohydrate reserves and sucrose in the medium to bulb growth of lily regenerated on scale segments *in vitro*. Physiologia Plantarum, 117: 245-255.

Larkindale J., Hall J. D., Knight M. R., Vierling E. (2005). Heat stress phenotypes of Arabidopsis mutants implicate multiple signaling pathways in the acquisition of thermotolerance. Plant Physiology, 138: 882-897.

Lian M. L., Chakrabarty D., Paek K. Y. (2003). Growth of *Lilium Oriental* Hybrid 'Casa blanca' bulblet using bioreactor culture. Scientia Horticulturae, 97: 41-48.

Murashige T., Skoog F. (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. Physiologia Plantarum, 15: 473-497.

Puijalon S., Piola F., Bornette G. (2008). Abiotic stresses increase plant regeneration ability. Evolutionary Ecology, 22: 493-506.

Pumisutapon P., Visser R. G. F., De Klerk G. J. (2012). Moderate abiotic stresses increase rhizome growth and outgrowth of axillary buds in Alstroemeria cultured *in vitro*. Plant Cell, Tissue and Organ Culture, 110: 395-400.

Qu L. W., Yin D. S., Su S. J., Pan B. T., Zhao X. H., Yang J. M., Pei X. H., Fu B. (2014). Production of lily (*Lilium*) bulbs and cut flowers in Liaoning Province of Northeast China. Acta Horticulturae, 1027: 105-112.

Robb S. M. (1957). The culture of excised tissue form bulb scales of *Lilium speciosum*Thun. Journal of Experimental Botany, 8: 348-352.

Takayama S., Misawa M. (1983). A scheme for mass propagation of *Lilium in vitro*. Scientia Horticulturae, 18: 353-362.

Tunnacliffe A., Wise M. J. (2007). The continuing conundrum of the LEA proteins. Naturwissenschaften, 94: 791-812.

Van Dam M. F. N. (2013). Controlling tulip stem nematodes in tulip bulbs by a hot water treatment. Acta Horticulturae, 1002: 327-331.

Wang W., Vinocur B., Altman A. (2003). Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. Planta, 218: 1-14.

Xin H., Zhang H., Chen L., Li X., Lian Q., Yuan X., Hu X., Cao L., He X., Yi M. (2010). Cloning and characterization of HsfA2 from Lily (*Lilium longiflorum*). Plant Cell Reports, 29: 875-885.