Propagation of Ornamental Plants
Vol. 2, № 1, 2002: 28-37

TRANSGENE EXPRESSION FOLLOWING TRANSFORMATION OF CHRYSANTHEMUM BY FOUR GENE INTRODUCTION METHODS

Jaime A.Teixeira da Silva and Seiichi Fukai

Faculty of Agriculture, Kagawa University, Miki-cho, Kagawa, 761-0795, Japan
E-mail: jaimetex@angelfire.com

REFERENCES

Benetka V., Pavingerová D. (1995). Phenotypic differences in transgenic plants of chrysanthemum. Plant Breeding, 114: 169-173.

Boase M. R., Miller R., Deroles S. C. (1997). Chrysanthemum systematics, genetics, and breeding. *In*: Janick J. (Ed.) Plant breeding reviews, vol. 14, John Wiley and Sons: 321-361.

Boase M. R., Bradley J. M., Borst N. K. (1998a). Genetic transformation mediated by *Agrobacterium tumefaciens* of florists' chrysanthemum (*Dendranthema* x *grandiflorum*) cultivar '*Peach Margaret*'. In Vitro Cellular and Developmental Biology - Plant, 34: 46-51.

Boase M. R., Butler R. C., Borst N. K. (1998 b). Chrysanthemum-cultivar-Agrobacterium interactions revealed by GUS expression time course experiments. Scientia Horticulturae, 77: 89-107.

Chasan R. (1994). Making sense (suppression) of viral RNA-mediated resistance. Plant Cell, 6: 1329-1331.

Chriqui D., David C., Adam S. (1988). Effect of the differentiated or dedifferentiated state of tobacco pith tissue on its behaviour after inoculation with *Agrobacterium rhizogenes*. Plant Cell Reports, 7: 111-114.

Courtney-Gutterson N., Napoli C., Lemieux C., Morgan A., Firoozabady E., Robinson K. E. P. (1994). Modification of flower color in florist's chrysanthemum: production of a white-flowering variety through molecular genetics. BioTechnology, 12: 268-271.

de Jong J., Rademaker W., van Wordragen M. F. (1993). Restoring adventitious shoot formation on chrysanthemum leaf explants following cocultivation with *Agrobacterium tumefaciens*. Plant Cell, Tissue and Organ Culture, 32: 263-270.

de Jong J., Mertens M. J., Rademaker W. (1994). Stable expression of the GUS reporter gene in chrysanthemum depends on binary plasmid T-DNA. Plant Cell Reports, 14: 59-64.

de Jong J., Rademaker W., Ohishi K. (1995). *Agrobacterium*-mediated transformation of chrysanthemum. Plant Tissue Culture and Biotechnology, 1: 38-42.

Dolgov S. V., Mitiouchkina T. Y., Skryabin K. G. (1997). Agrobacterial transformation of chrysanthemum. Acta Horticulturae, 447: 329-333.

Fu R-Z., Liu M., Liang H-J., Zhang C-H., Xue H., Sun Y-R. (1998). Production of transgenic plants of chrysanthemum via *Agrobacterium tumefaciens* mediated method. Acta Phytophysiologica Sinica, 24: 72-76.

Fukai S., Chen Z., Oe M. (1987). Cultivar differences in adventitious shoot formation from leaf segments of chrysanthemum *(Dendranthema grandiflora* (Ramat.) Kitamura). Bulletin of the Osaka Agricultural Research Centre, 24: 55-58.

Fukai S., de Jong J., Rademaker W. (1995). Efficient genetic transformation of chrysanthemum (*Dendranthema grandiflora* (Ramat.) Kitamura) using stem segments. Breeding Science, 45: 179-184.

Horsch R. B., Fry J. E., Hoffman N. L., Eichholtz D., Rogers S. G., Fraley R. T. (1985). A simple and general method for transferring genes into plants. Science, 227: 1229-1231.

Kamo K., Blowers A. (1999). Tissue specificity and expression level of *gus*A under *rol*D, mannopine synthase and translation elongation factor 1 subunit α promoters in transgenic *Gladiolus* plants. Plant Cell Reports, 18: 809-815.

Jefferson R. A., Kavanagh T. A., Bevan M. W. (1987). GUS fusions: β-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. EMBO Journal, 6: 3901-3907.

Ledger S. E., Deroles S. C., Given N. K. (1991). Regeneration and *Agrobacterium*-mediated transformation of chrysanthemum. Plant Cell Reports, 10: 195-199.

Lowe J. M., Davey M. R., Power J. B., Blundy K. S. (1993). A study of some factors affecting *Agrobacterium*-transformation and plant regeneration of *Dendranthema grandiflora* Tzvelev (syn. *Chrysanthemum morifolium* Ramat.). Plant Cell, Tissue and Organ Culture, 33: 171-180.

Mishiba K., Mii K. (2000). Polysomaty analysis in diploid and tetraploid *Portulaca grandiflora*. Plant Science 156: 213-219.

Murray M. G., Thompson W. F. (1980). Rapid isolation of high molecular weight plant DNA. Nucleic Acids Research, 8: 4321-4325.

Odell J. T., Nagy F., Chua N-H. (1985). Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. Nature, 313: 810-812.

Pavingerová D., Dostál J., Bisková R., Benetka V. (1994). Somatic embryogenesis and *Agrobacterium*-mediated transformation of chrysanthemum. Plant Science, 97: 95-101.

Renou J. P., Brochard P., Jalouzot R. (1993). Recovery of transgenic chrysanthemum (*Dendranthema grandiflora* Tzvelev) after hygromycin resistance selection. Plant Science, 89: 185-197.

Sherman J. M., Moyer J. W., Daub M. E. (1998). A regeneration and *Agrobacterium*-mediated transformation system for genetically diverse chrysanthemum cultivars. Journal of the American Society for Horticultural Science, 123: 189-194.

Shinoyama H., Komano M., Nomura Y., Kazuma T. (1998). Stable *Agrobacterium*-mediated transformation of chrysanthemum (*Dendranthema grandiflora* (Ramat.) Kitamura). Bulletin of the Fukui Agricultural Experimental Station, 35: 13-21.

Takatsu Y., Nishizawa Y., Hibi T., Akutsu K. (1999). Transgenic chrysanthemum (*Dendranthema grandiflora* (Ramat.) Kitamura) expressing a rice chitinase gene shows enhanced resistance to gray mold (*Botrytis cinerea*). Scientia Horticulturae, 82: 113-123.

Tosca A., Delledonne M., Furini A., Belenghi B., Fogher C., Frangi P. (2000). Transformation of Korean chrysanthemum (*Dendranthema zawadskii x D.* x *grandiflorum*) and insertion of the maize autonomous element Ac using *Agrobacterium tumefaciens*. Journal of Genetics and Breeding, 54: 19-24.

Urban L. A., Sherman J. M., Moyer J. W., Daub M. E. (1994). High frequency shoot regeneration and *Agrobacterium*-mediated transformation of chrysanthemum (*Dendranthema grandiflora*). Plant Science, 98: 69-79.

Vancanneyt G., Schmidt R., O'Connor-Sanchez A., Willmitzer L., Rocha-Sosa M. (1990). Construction of an intron-containing marker gene: splicing of the intron in transgenic plants and its use in monitoring early events in *Agrobacterium*-mediated plant transformation. Molecular and General Genetics, 220: 245-50.

van Wordragen M. F., de Jong J., Huitema H. B. M., Dons H. J. M. (1991). Genetic transformation of chrysanthemum using wild type *Agrobacterium* strains; strain and cultivar specificity. Plant Cell Reports, 9: 505-508.

van Wordragen M. F., Ouwerkerk P. B. F., Dons H. J. M. (1992 a). *Agrobacterium rhizogenes* mediated induction of apparently untransformed roots and callus in chrysanthemum. Plant Cell, Tissue and Organ Culture, 30: 149-157.

van Wordragen M. F., De Jong J., Schornagel M. J., Dons H. J. M. (1992 b). Rapid screening for host-bacterium interactions in *Agrobacterium*-mediated gene transfer to chrysanthemum, by using the GUS-intron gene. Plant Science, 81: 207-214.

Yepes L. C., Mittak V., Pang S-Z., Gonsalves C., Slightom J. L., Gonsalves D. (1995). Biolistic transformation of chrysanthemum with the nucleocapsid gene of tomato spotted wilt virus. Plant Cell Reports, 14: 694-698.

Young K. J., Jung P. S., Young U. B., Ho P. C., Soo C. Y., Sheop S. J. (1998). Transformation of chrysanthemum by *Agrobacterium tumefaciens* with three different types of vectors. Journal of the Korean Society for Horticultural Science, 39: 360-366.

Zheng Z-L., Yang Z., Jang J-C., Metzger J. D. (2001). Modification of plant architecture in chrysanthemum by ectopic expression of the tobacco phytochrome *B1* gene. Journal of the American Society for Horticultural Science, 126: 19-26.