

***CURCUMA*: STUDIES ON TISSUE CULTURE, POLLEN GERMINATION AND VIABILITY,
HISTOLOGY AND FLOW CYTOMETRY**

Waraporn Udomdee¹, Seiichi Fukai², Luckana Petpradap¹ and Jaime A. Teixeira da Silva^{2*}

¹Department of Horticulture, Maejo University, Chiang Mai, 50290, Thailand

²Faculty of Agriculture, Kagawa University, Miki-cho, Ikenobe, 2393, Kagawa-ken, 761-0795, Japan,
Telfax: +81 (0)87 891 0747, E-mail: jaimetex@angelfire.com

REFERENCES

- Adaniya S. (2001). Optimal pollination environment of tetraploid ginger (*Zingiber mioga* Roscoe) evaluated by *in vitro* pollen germination and pollen tube growth in styles. *Scientia Horticulturae*, 90: 219-226.
- Adaniya S., Shirai D. (2001). *In vitro* induction of tetraploid ginger (*Zingiber officinale* Roscoe) and its pollen fertility and germinability. *Scientia Horticulturae*, 88: 277-287.
- Apavatjirut P., Sirisawad T., Sirirugsa P., Voraurai P., Suwanthada C. (1996). Studies on chromosome number of seventeen Thai *Curcuma* species. Proceedings 2nd national conference on flowers and ornamentals, Thailand, 2: 86-93. (In Thai with English abstract).
- Banjerpongchai R., Suttajit M., Kongtawelert P. (2002). Effect of curcumin on HL60 cell apoptosis induced by doxorubicin. *Chiang Mai Medical Bulletin*, 41: 59-66.
- Barre P., Noirot M., Louarn J., Duperray C., Hamon S. (1996). Reliable flow cytometric estimation of nuclear DNA content in coffee trees. *Cytometry*, 24: 32-38.
- Bennett M. D., Leitch I. J. (1995). Nuclear DNA amounts in angiosperms. *Annals of Botany*, 76: 113-176.
- Bowes S. A. (1990). Long-term storage of *Narcissus* anthers and pollen in liquid nitrogen. *Euphytica*, 48: 275-278.
- Chong-Jin G., Nathan M. J., Kumar P. P. (1995). Direct organogenesis and induction of morphogenic callus through thin section culture of *Heliconia psittacorum*. *Scientia Horticulturae*, 62: 113-120.
- De Laat A. N. M., Göhde W., Vogelzang M. J. D. (1987). Determination of ploidy of single plants and plant populations by flow cytometry. *Plant Breeding*, 99: 303-307.
- Faria R. T., Illg R. D. (1995). Micropropagation of *Zingiber spectabile* Griff. *Scientia Horticulturae*, 62: 135-137.
- Hagiladi A., Umiel N., Gilad Z., Yang X. H. (1997). *Curcuma alismatifolia*. I. Plant morphology and effect of tuberous root number on flowering date and yield of inflorescences. *Acta Horticulturae*, 430: 747-753.
- Hughes H. G., Lee C. W., Towill L. E. (1991). Low-temperature preservation of *Clanthus formosus* pollen. *HortScience*, 26: 1411-1412.
- Ikeda H., Numata S. (1998). Pollen storage of chrysanthemum. *Acta Horticulturae*, 454: 329-333.
- Iwanami Y. (1972). Retaining the viability of *Camellia japonica* pollen in various organic solvents. *Plant and Cell Physiology*, 13: 1139-1141.
- Meenakshi N., Sulikeri G.S., Krishnamoorthy V., Hegde R.V. (2001). Standardisation of chemical environment for multiple shoot induction of turmeric (*Curcuma longa* L.) for *in vitro* clonal propagation. *Crop Research*, 22: 449-453.
- Mishiba K., Ando T., Mii M., Watanabe H., Kokubun H., Hashimoto G., Marchesi E. (2000). Nuclear DNA content as an index character discriminating taxa in the genus *Petunia sensu* Jussieu (*Solanaceae*). *Annals of Botany*, 85: 665-673.
- Mukhri Z., Yamaguchi H. (1986). *In vitro* plant multiplication from rhizomes of turmeric (*Curcuma domestica* val) and Temoe Lawak (*C. xanthorrhiza* Robx.). *Plant Tissue Culture Letters*, 3: 28-30.
- Murashige T., Skoog F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- Nayak S. (2000). *In vitro* multiplication and microrhizome induction in *Curcuma aromatica* Salisb. *Plant Growth Regulation*, 32: 41-47.

- Niimi Y., Shiokawa Y. (1992). A study on the storage of *Lilium* pollen. Journal of the Japanese Society for Horticultural Science, 61: 399-403.
- Panyabal (2000). Tissue culture of *Curcuma* sp. <http://www.rb.ac.th/org/research/rajabhat/rilp/cs3.html>.
- Patyanun P. (2000). The appropriate position on *Curcuma* inflorescences as initial explant for micropropagation scheme. Faculty of Science, Maejo University, Thailand, pp. 21 (In Thai with English abstract).
- Pubuopiend J. (1997). Growth and floral development of *Curcuma sparganifolia* Gagnep. H. MSc. Thesis, Chiang Mai University (In Thai with English abstract).
- Rajasekharan P. E., Rao T. M., Janakiram T., Ganeshan S. (1994). Freeze preservation of gladiolus pollen. Euphytica, 80: 105-109.
- Salvi N. D., George L., Eapen S. (2001). Plant regeneration from leaf base callus of turmeric and random amplified polymorphic DNA analysis of regenerated plants. Plant Cell, Tissue and Organ Culture, 66: 113-119.
- Sharma T. R., Singh B. M. (1997). High-frequency *in vitro* multiplication of disease-free *Zingiber officinale* Rosc. Plant Cell Reports, 17: 68-72.
- Wannakrairoj S. (1992). *In vitro* propagation of Patumma (*Curcuma alismatifolia* Gagnep.). HortScience, 27: 97.
- Wannakrairoj S. (1997). Curcuma. Amarin Printing and Publishing Public Co., Ltd, 128 pp. (In Thai).
- Yokoya K., Roberts A. V., Mottley J., Lewis R., Brandham P. E. (2000). Nuclear DNA amounts in roses. Annals of Botany, 85: 557-561.
- Yu-Mei H. (1996). Effects of corm size and number of storage roots on flowering date and yield of flower stalks and corms of *Curcuma alismatifolia*. Research Bulletin of Kaohsiung District Agricultural Improvement Station, Taiwan, 7: 13-17 (In Chinese with English abstract).