

TISSUE CULTURE OF CHRYSANTHEMUM: A REVIEW

Jaime A. Teixeira da Silva

Faculty of Agriculture, Kagawa University, Miki-cho, Ikenobe, 2393, Kagawa-ken, 761-0795, Japan
Telfax: +81 (0) 87 891 0747, E-mail: jaimetex@lycos.com

REFERENCES

- Ahmed H. A. (1986). *In vitro* regeneration and propagation of meristem apices of chrysanthemum. Kerteszet-egyetem Közleményei, 50: 199-214.
- Ahmed H. A., Andrea M. (1987). Effect of *Chrysanthemum* multiplication by meristem-tip culture. Acta Horticulturae, 212: 98-99.
- Amagasa K., Kameya T. (1989). Plant regeneration and callus formation from *Chrysanthemum morifolium*, *C. coronarium* and *Lactuca sativa* protoplasts. Journal of the Japanese Society for Horticultural Science, 57: 620-625.
- Anderson N. O., Ascher P. D., Widmer R. E., Luby J. J. (1990). Rapid regeneration cycling of chrysanthemum using laboratory seed development and embryo rescue techniques. Journal of the American Society for Horticultural Science, 115: 329-336.
- Annadana S., Rademaker W., Ramanna M., Udayakumar M., de Jong J. (2000). Response of stem explants to screening and explant source as a basis for methodological advancing of regeneration protocols for chrysanthemum. Plant Cell, Tissue and Organ Culture, 62: 47-55.
- Annadana S., Mlynárová L., Udayakumar M., de Jong J., Nap J. P. (2001). The potato *Lhca3.St.1* promoter confers high and stable transgene expression in chrysanthemum, in contrast to CaMV-based promoters. Molecular Breeding, 8: 335-344.
- Aribaud M., Kevers C., Martin-Tanguy J., Gaspar T. (1999). Low activity of amine-oxidases and accumulation of conjugated polyamines in disfavour of organogenic programs in chrysanthemum leaf disk explants. Plant Cell, Tissue and Organ Culture, 55: 85-94.
- Bajaj Y. P. S., Sidhu M. M. S., Gill A. P. S. (1992). Micropropagation of *Chrysanthemum*. In: Bajaj Y. P. S. (Ed.). Biotechnology in Agriculture and Forestry, Vol. 20: High-tech and micropropagation IV. Springer-Verlag, Berlin, Germany: 69-80.
- Bannier L. J., Steponkus P. L. (1976). Cold acclimation of chrysanthemum callus cultures. Journal of the American Society for Horticultural Science, 101: 409-412.
- Barthomeuf C., Hitmi A., Veisseire P., Coudret A. (1996). Identification and assay of pyrethrins in *Chrysanthemum cinerariaefolium* calli. Biotechnology Techniques, 10: 639-642.
- Benetka V., Pavingerová D. (1995). Phenotypic differences in transgenic plants of chrysanthemum. Plant Breeding, 114: 169-173.
- Ben-Jaacov J., Langhans R. W. (1972). Rapid multiplication of *Chrysanthemum* plants by stem tip proliferation. HortScience, 7: 289-290.
- Bhattacharya P., Dey S., Das N., Bhattacharya B. C. (1990). Rapid mass propagation of *Chrysanthemum morifolium* by callus derived from stem and leaf explants. Plant Cell Reports, 9: 439-442.
- Bhattacharya P., Dey S., Bhattacharya B. C. (1994). Use of low-cost gelling agents and support matrices for industrial scale plant tissue culture. Plant Cell, Tissue and Organ Culture, 37: 15-23.
- Boase M. R., Bradley J. M., Borst N. K. (1998a). Genetic transformation mediated by *Agrobacterium tumefaciens* of florists' chrysanthemum (*Dendranthema x grandiflorum*) cultivar 'Peach Margaret'. In Vitro Cellular and Developmental Biology – Plant, 34: 46-51.
- Boase M. R., Borst N. K., Bradley J. M. (1998b). Chrysanthemum cultivar-*Agrobacterium* interactions revealed by GUS expression time course experiments. Scientia Horticulturae, 77: 89-107.
- Broertjies C., Roest S., Bokelmann G. S. (1976). Mutation breeding of *Chrysanthemum morifolium* Ram. using *in vivo* and *in vitro* adventitious bud techniques. Euphytica, 25: 11-19.
- Bush R., Earle E. D., Langhans R. W. (1976). Plantlets from petal segments, petal epidermis and shoot tips of the periclinal chimera, *Chrysanthemum morifolium* 'Indianapolis'. American Journal of Botany, 63: 729-737.
- Chakrabarty D., Mandal A. K. A., Datta S. K. (1999). Management of chimera through direct shoot regeneration from florets of chrysanthemum (*Dendranthema grandiflora* Ramat.). Journal of Horticultural Science and

- Biotechnology, 74: 293-296.
- Chakrabarty D., Mandal A. K. A., Datta S. K. (2000). SEM and light microscopic studies on direct shoot regeneration from ray florets of *Chrysanthemum*. Israel Journal of Plant Sciences, 48: 105-107.
- Chen Y. Z., He X. D., Jiang P. Y., Wang C. M. (1985). *In vitro* propagation of *Chrysanthemum* leaves. Journal of the Jiangsu Agricultural College, 6: 33-36.
- Chen Z. L., Hou S. W., Yu H. Y. (1998). Tissue culture of *Pyrethrum cinerariaefolium*. Acta Botanica Yunnanica, 20: 351-354.
- Chumsri P., Staba E. J. (1975). Pyrethrins: content and larvicidal activity of *Chrysanthemum* plants and tissue cultures. Academia Pharmacia Scientiae, Abstracts 5: 169
- Chung B. N., Choi G. S., Kim H. R., Kim J. S. (2001) Chrysanthemum stunt viroid in *Dendranthema grandiflorum*. Plant Pathology Journal, 17: 194-200.
- Corneanu G. C., Corneanu M. (1992). Preliminary studies about human bio-energy effect on *in vitro* vegetal cultures. Révue Roumaine de Biologie, 37: 113-117.
- Courtney-Gutterson N., Napoli C., Lemieux C., Morgan A., Firoozabady E., Robinson K. E. P. (1994). Modification of lower color in florist's chrysanthemum: production of a white-flowering variety through molecular genetics. BioTechnology, 12: 268-271.
- Courtney-Gutterson N., Otten A., Firoozabady E., Akerboom M., Lemieux C., Nicholas J., Morgan A., Robinson K. E. P. (1993). Production of genetically engineered color-modified chrysanthemum plants carrying a homologous chalcone synthase gene and their field performance. Acta Horticulturae, 336: 57-62.
- Dabin P., Choiseg D. (1983). Application of *in vitro* culture to the vegetative propagation of *Chrysanthemum* cv. *White Spider*. Bulletin de la Société Royale de Botanique Belgique, 116: 161-166.
- Datta S. K., Chakrabarty D., Saxena M., Mandal A. K. A., Biswas A. K. (2001). Direct shoot regeneration from florets of chrysanthemum cultivars. Indian Journal of Genetics, 61: 373-376.
- Daughtrey M. L., Jones R. K., Moyer J. W., Daub M. E., Baker J. R. (1997). Tospoviruses strike the greenhouse industry: INSV has become a major pathogen on flower crops. Plant Disease, 81: 1220-1230.
- de Argollo M. D., Kirszenzaft S. S. L., Jesu C. O. (1998). Influence of nitrogen sources on *in vitro* morphogenesis of axillary shoots in stem explants of *Chrysanthemum morifolium* Ramat. Revista Brasileira de Botânica, 21: 141-147.
- de Donato M., Peruceo E. (1984). Micropropagation of *Chrysanthemum* by means of lateral meristem stimulation. Annali di Faculti Scientia Agraria de la Universitati Studi Torino, 13: 103-115.
- de Jong J., Custers J. B. M. (1986). Induced changes in growth and flowering of chrysanthemum after irradiation and *in vitro* culture of pedicels and petal epidermis. Euphytica, 35: 137-148.
- de Jong J., Van Wordragen M. F., Rademaker W. (1990). Early transformation events in *Dendranthema grandiflora*. In: Integration of *in vitro* techniques in ornamental plant breeding. Proceedings of the EUCARPIA (Section Ornamentals), Wageningen: 156-161.
- de Jong J., Rademaker W., van Wordragen M. F. (1993). Restoring adventitious shoot formation on chrysanthemum leaf explants following cocultivation with *Agrobacterium tumefaciens*. Plant Cell, Tissue and Organ Culture, 32: 263-270.
- de Jong J., Mertens M. J., Rademaker W. (1994). Stable expression of the GUS reporter gene in chrysanthemum depends on binary plasmid T-DNA. Plant Cell Reports 14: 59-64.
- de Jong J., Rademaker W., Ohishi K. (1995). *Agrobacterium*-mediated transformation of chrysanthemum. Plant Tissue Culture and Biotechnology, 1: 38-42.
- de Oliveira P. D., Moacir P., Renato P. (1995). The effect of different concentrations of growth regulators on *in vitro* shoot proliferation of chrysanthemum (*Dendranthema grandiflora* Tzvelev). Ciência e Prática, 19: 397-408.
- de Oliveira P. D., Pasqual M., Paiva R. (1996). Different concentrations of the MS medium, nitrogen and sucrose on micropropagation of chrysanthemum. Bragantia, 55: 9-18.
- Dhar K., Pal A. (1993). Factors influencing efficient pyrethrin production in undifferentiated cultures of *Chrysanthemum cinerariaefolium*. Fitoterapia, 64: 336-340.
- Dolgov S. V., Mityshkina T. U., Rukavtsova E. B., Buryanov Y. I. (1995). Production of transgenic plants of *Chrysanthemum morifolium* Ramat. with the gene of *Bac. thuringiensis* δ -endotoxin. Acta Horticulturae, 441: 223-228.
- Dolgov S. V., Mitiouchkina T. Y., Skryabin K. G. (1997). Agrobacterial transformation of chrysanthemum. Acta Horticulturae, 447: 329-333.
- Dwivedi A. K., Banerji B. K., Chakrabarty D., Mandal A. K. A., Datta S. K. (2000). Gamma ray induced new flower colour chimera and its management through tissue culture. Indian Journal of Agricultural Science, 70: 853-855.
- Earle E. D., Langhans R. W. (1974a). Propagation of chrysanthemum *in vitro* I. Multiple plantlets from shoot

- tips and the establishment of tissue cultures. *Journal of the American Society for Horticultural Science*, 99: 128-131.
- Earle E. D., Langhans R. W. (1974b). Propagation of chrysanthemum *in vitro* II. Production growth and flowering of plantlets from tissue cultures. *Journal of the American Society for Horticultural Science*, 99: 352-358.
- Endo M., Sasaki T., Inada I. (1990). Creation of mutants through tissue culture of edible chrysanthemums *Chrysanthemum morifolium* Ram. I. Especially the relationship among the different explants and variation in their regenerated plants. *Journal of the Faculty of Agriculture Iwate University*, 20: 17-34.
- Endo M., Sasaki T., Inada I. (1991). Creation of mutants through tissue culture of edible chrysanthemums *Chrysanthemum morifolium* Ram. II. Especially an attempt to create chromosome-reduced plants by treatment with PFP on agar medium. *Journal of the Faculty of Agriculture Iwate University*, 20: 91-103.
- Endo M., Inada I. (1997). Production and characteristics of chromosome-doubled plants of small-flowered garden chrysanthemum, *Dendranthema grandiflorum* (Ramatt.) Kitam. cv. *YS* by colchicines treatment of cultured shoot tips. *Journal of the Japanese Society for Horticultural Science*, 65: 825-833.
- Flowers and Plants Association. <http://flowers.org.uk/flowers/trivia/toptenflowers2.htm>. 2001.
- Fu R. Z., Liu M., Liang H. J., Zhang C. H., Xue H., Sun Y. R. (1998). Production of transgenic plants of chrysanthemum via *Agrobacterium tumefaciens* mediated method. *Acta Phytophysiologica Sinica*, 24: 72-76.
- Fujii Y., Shimizu K. (1990). Regeneration of plants from achenes and petals of *Chrysanthemum coccineum*. *Plant Cell Reports*, 8: 625-627.
- Fukai S. (1986). Effects of sugar on callus and organ formation from leaf segments of chrysanthemum *Dendranthema grandiflorum* Kitamura). *Bulletin of the Osaka Agricultural Research Center*, 3: 71-77.
- Fukai S. (1990). Cryopreservation of chrysanthemum shoot tips. *Scientia Horticulturae*, 45: 167-174.
- Fukai S. (1995). Cryopreservation of germplasm of chrysanthemums. In: Bajaj Y. P. S. (Ed.). *Biotechnology in Agriculture and Forestry*, Vol. 32. *Cryopreservation of Plant Germplasm I*. Springer-Verlag Berlin: 447-457.
- Fukai S., Oë M. (1986). Effects of plant growth regulators on organ formation from leaf and stem segments of chrysanthemum. *Bulletin of the Osaka Agricultural Research Center*, 23: 25-31.
- Fukai S., Oë M. (1990). Morphological observations of chrysanthemum shoot tips cultured after cryoprotection and freezing. *Journal of the Japanese Society for Horticultural Science*, 59: 383-387.
- Fukai S., Goi M., Tanaka M. (1994). The chimeric structure of the apical dome of chrysanthemum (*Dendranthema grandiflorum* (Ramat. Kitam.) is affected by cryopreservation. *Scientia Horticulturae*, 57: 347-351.
- Fukai S., de Jong J., Rademaker W. (1995). Efficient genetic transformation of chrysanthemum (*Dendranthema grandiflorum* (Ramat.) Kitamura) using stem segments. *Breeding Science*, 45: 179-184.
- Gao Y., Zhao B., Ding G., Zhang Q. (2001). Shoot regeneration from stem and leaf explants of *Dendranthema grandiflorum*. *Journal of the Beijing Forestry University*, 23: 32-33.
- George J., Rajasekaran T., Ravishankar G. A. (1999). A modified culture vessel for improved callus growth and pyrethrins content of pyrethrum (*Chrysanthemum cinerariaefolium* Vis.). *Pyrethrum Post*, 20: 49-54.
- George M. W., Tripepi R. R. (2001). Plant Preservative Mixture® can affect shoot regeneration from leaf explants of chrysanthemum, European birch, and rhododendron. *HortScience*, 36: 768-769.
- Gertsson U. E., Andersson E. (1985). Propagation of *Chrysanthemum x hortorum* and *Philodendron scandens* by tissue culture. *Rapport Institute Tradgards Svërig Landbruks University*, 41: 17.
- Grewal S., Sharma K. (1978). Pyrethrum plant (*Chrysanthemum cinerariaefolium* Vis.) regeneration from shoot tip culture. *Indian Journal of Experimental Biology*, 16: 1119-1121.
- Hahn E., Bae J., Lee J. H., Beom Y. (1998). Growth and leaf-surface characteristics of chrysanthemum plantlets between hydroponic and microponic system. *Journal of the Korean Society for Horticultural Science*, 39: 838-842.
- Harborne J. B., Heywood V. H., Saleh N. A. M. (1970). Chemosystematics of the Compositae: flavonoid patterns in the *Chrysanthemum* complex of the tribe Anthemideae. *Phytochemistry*, 9: 2011-2017.
- Hattori K. (1992). The process during shoot regeneration in the receptacle culture of chrysanthemum, *Chrysanthemum morifolium* Ramat. *Japanese Journal of Breeding*, 42: 227-234.
- Hill G. P. (1968). Shoot formation in tissue cultures of chrysanthemum 'Bronze Pride'. *Physiologia Plantarum*, 2: 386-389.
- Hitmi A., Barthomeuf C., Sallanon H. (2000). Cryopreservation of *Chrysanthemum cinerariaefolium* shoot tips. *Journal of Plant Physiology*, 156: 408-412.
- Iizuka M., Matsumoto E., Doi A., Madrigal R., Fukushima A. (1973). Tubular floret culture of chrysanthemum and cineraria *in vitro*. *Japanese Journal of Genetics*, 48: 79-87.
- Jeong J. H., Chakrabarty D., Kim S. J., Paek K. Y. (2002). Transformation of chrysanthemum (*Dendranthema grandiflorum* Kitamura cv. *Cheonsu* by constitutive expression of rice *OsMADS1* gene. *Journal of the Korean Society for Horticultural Science*, 43: 382-386.

- Kaul V., Miller R. M., Hutchinson J. F., Richards D. (1990). Shoot regeneration from stem and leaf explants of *Dendranthema grandiflora* Tzvelev (syn. *Chrysanthemum morifolium* Ramat.). *Plant Cell, Tissue and Organ Culture*, 21: 21-30.
- Kaur-Sawhney R., Applewhite P. B. (1993). Endogenous protein-bound polyamines: correlation with regions of cell division in tobacco leaves, internodes and ovaries. *Plant Growth Regulation*, 12: 223-227.
- Khalid N., Davey M. R., Power J. B. (1989). An assessment of somaclonal variation in *Chrysanthemum morifolium*: the generation of plants of potential commercial value. *Scientia Horticulturae*, 38: 287-294.
- Khehra M., Lowe K. C., Davey M. R., Power J. B. (1995). An improved micropropagation system for *Chrysanthemum* based on Pluronic F-68-supplemented media. *Plant Cell, Tissue and Organ Culture*, 41: 87-90.
- Kim J., Park Y., Jung S., Chung H., Shin Y. S., Sheop J. (1998). Transformation of chrysanthemum by *Agrobacterium tumefaciens* with three vectors. *Journal of the Korean Society for Horticultural Science*, 39: 360-366.
- Kim M., Kim J., Hee Y. (1998). Plant regeneration and flavonoid 3',5'-hydroxylase gene transformation of *Dendranthema indicum* and *Dendranthema zawadskii*. *Journal of the Korean Society for Horticultural Science*, 39: 355-359.
- Kohno H., Ishigeki M., Yoshida F. (1978). Comparative studies of mineral nutrition calli induced from various plant species. I. Callus induction from various plant species. *Bulletin of the Faculty of Agriculture Tamagawa University*, 18: 16-21.
- Krishna D., Pal A., Dhar K., Pal A. (1993). Factors influencing efficient pyrethrin production in undifferentiated cultures of *Chrysanthemum cinerariaefolium*. *Fitoterapia*, 64: 336-340.
- Kudo S., Shibata N., Kanno Y., Suzuki M. (2002). Transformation of chrysanthemum (*Dendranthema grandiflorum* (Ramat.) Kitamura) via *Agrobacterium tumefaciens*. *Acta Horticulturae*, 572: 139-147.
- Kumar A., Kumar V. A. (1995). High-frequency *in vitro* propagation in *Chrysanthemum maseimum*. *Indian Horticulture*, 1: 37-38.
- Lazar M., Cachita C. D. (1982). Micropropagation of Chrysanthemums. II. *In vitro* culture of shoot meristems. *Product and Vegetable Horticulture*, 31: 32-35.
- Lazar M., Cachita C. D. (1983). Micropropagation of Chrysanthemus. III. Chrysanthemum multiplication *in vitro* from capitulum explants. *Product and Vegetable Horticulture*, 32: 44-47.
- Ledger S. E., Derolles S. C., Given N. K. (1991). Regeneration and *Agrobacterium*-mediated transformation of chrysanthemum. *Plant Cell Reports*, 10: 195-199.
- Lee J. K., Park K. Y., Chun C. K. (1979). *In vitro* propagation of *Chrysanthemum* through shoot apical meristem culture. *Journal of the Korean Society for Horticultural Science*, 20: 192-199.
- Lee T., Huang M. E. E., Pua E. C. (1997). High frequency shoot regeneration from leaf disk explants of garland chrysanthemum (*Chrysanthemum coronarium* L.) *in vitro*. *Plant Science*, 126: 219-226.
- Lee Y., Kwon K., Lee Y. J. (1999). Plant regeneration from leaf segment cultures of chrysanthemum (*Dendranthema grandiflora* Tzvelev). *Korean Journal of Plant Tissue Culture*, 26: 59-63.
- Lemieux C. S., Firoozabady E., Robinson K. E. P. (1990). *Agrobacterium*-mediated transformation of chrysanthemum. In: de Jong J. (Ed.). *Integration of in vitro techniques in ornamental plant breeding*. Wageningen, CPO Press: 150-155.
- Lindsay G. C., Ledger S. E. (1993). A protoplast to plant system for chrysanthemum *Dendranthema zawadskii* x *D. grandiflora*. *Plant Cell Reports*, 12: 278-280.
- Lowe J. M., Davey M. R., Power J. B., Blundy K. S. (1993). A study of some factors affecting *Agrobacterium*-transformation and plant regeneration of *Dendranthema grandiflora* Tzvelev (syn. *Chrysanthemum morifolium* Ramat.). *Plant Cell, Tissue and Organ Culture*, 33: 171-180.
- Lu C. Y., Nugent G., Wardley T. (1990). Efficient direct plant regeneration from stem segments of chrysanthemum (*Dendranthema morifolium* Ramat. cv. Royal Purple). *Plant Cell Reports*, 8: 733-736.
- Malaure R. S., Davey M. R., Power J. B. (1989). Isolation and culture of protoplasts of *Chrysanthemum cinerariaefolium* Vis. *Pyrethrum Post*, 17: 90-94.
- Malaure R. S., Barclay G., Power J. B., Davey M. R. (1991a). The production of novel plants from florets of *Chrysanthemum morifolium* using tissue culture. 1. Shoot regeneration from ray florets and somaclonal variation exhibited by regenerated plants. *Journal of Plant Physiology*, 139: 8-13.
- Malaure R. S., Barclay G., Power J. B., Davey M. R. (1991b). The production of novel plants from florets of *Chrysanthemum morifolium* using tissue culture. 2. Securing natural mutations (sports). *Journal of Plant Physiology*, 139: 14-18.
- Mandal A. K. A., Chakrabarty D., Datta S. K. (2000a). *In vitro* isolation of solid novel flower colour mutants from induced chimeric ray florets of chrysanthemum. *Euphytica*, 114: 9-12.
- Mandal A. K. A., Chakrabarty D., Datta S. K. (2000b). Application of *in vitro* techniques in mutation breeding of chrysanthemum. *Plant Cell, Tissue and Organ Culture*, 60: 33-38.

- May R. A., Trigiano R. N. (1991). Somatic embryogenesis and plant regeneration from leaves of *Dendranthema grandiflora*. *Journal of the American Society for Horticultural Science*, 116: 366-371.
- Mitra A., Bhattacharya P. S., Dey S., Sawarkar S. K., Bhattacharya B. C. (1998). Photoautotrophic *in vitro* culture of chrysanthemum under CO₂ enrichment. *Biotechnology Techniques*, 12: 335-337.
- Miyazaki S., Tashiro Y., Shimada T. (1976). Tissue culture of *Chrysanthemum morifolium* Ramat. I. Cultivar differences in organ formation. *Agricultural Bulletin of Saga University*, 40: 31-44.
- Miyazaki S., Tashiro Y. (1978). Tissue culture of *Chrysanthemum morifolium* Ramat. IV. On the explant sources for stem segment culture. *Agricultural Bulletin of Saga University*, 44: 67-78.
- Miyazaki S., Kishida E., Tashiro Y., Kanazawa K. (1979). Tissue culture of *Chrysanthemum morifolium* Ramat. V. Histological studies on the callus and shoot formation in stem segments cultured *in vitro*. *Agricultural Bulletin Saga University*, 46: 43-65.
- Mizutani T., Tanaka T. (1994). Study on the floret culture of Higo-chrysanthemum. *Proceedings of the School of Agriculture, Kyushu Tokai University*, 13: 9-14.
- Nhut D. T., Teixeira da Silva J. A., Aswath C. R. (2003). The importance of the explant on regeneration in thin cell layer technology. *In Vitro Cellular and Developmental Biology*, 39: 266-276.
- Ohishi K., Sakurai Y. (1988). Morphological changes in Chrysanthemum derived from petal tissue. *Research Bulletin of the Aichiken Agricultural Research Center*, 20: 278-284.
- Oka S., Muraoka O., Abe T., Nakajima S. (1996). Formation of leaf-like bodies and adventitious buds, and chimeric expression of introduced GUS gene in garland chrysanthemum tissue cultures. *Journal of the Japanese Society for Horticultural Science*, 65: 294-295.
- Oka S., Muraoka O., Abe T., Nakajima S. (1999). Adventitious bud and embryoid formation in garland chrysanthemum leaf culture. *Journal of the Japanese Society for Horticultural Science*, 68: 70-72.
- Okamura M., Hayashi T., Miyazaki S. (1984). Inhibiting effect of ammonium ion in protoplast culture of some Asteraceae plants. *Plant Cell Physiology*, 25: 281-286.
- Otsuka H., Suematsu N., Toda M. (1985). The culture and plant regeneration from mesophyll protoplast of chrysanthemum. *Bulletin of the Shizuoka Agricultural Experimental Station*, 30: 25-33.
- Otsuka H., Yamada H., Suematsu N., Toda M. (1987). Somaclonal variation of protoplast-derived regenerates in chrysanthemum. *Bulletin of the Shizuoka Agricultural Experimental Station*, 32: 53-59.
- Panfilová O. F., Andrianov V. N. (1996). Electrostimulation of rooting of chrysanthemum green cuttings after long-term storage. *Izvestiya Timiryazevskoi Selskokozyaistvennoi Akademii* 10: 105-113.
- Paul A., Dhar K., Pal A. (1988). Organogenesis from selected culture lines of pyrethrum *Chrysanthemum cinerariaefolium*. *Pyrethrum Post*, 17: 17-20.
- Pavingerová D., Dostál J., Bisková R., Benetka V. (1994). Somatic embryogenesis and *Agrobacterium*-mediated transformation of chrysanthemum. *Plant Science*, 97: 95-101.
- Pillai V., Zulkifli L. (2000). Somaclonal variation in *Chrysanthemum morifolium* generated through petal cultures. *Journal of Tropical Agriculture and Food Science*, 28: 115-120.
- Prasad R. N., Sharma A. K., Chaturvedi H. C. (1983). Clonal multiplication of *Chrysanthemum morifolium* 'Otome zakura' in long-term culture. *Bangladesh Journal of Botany*, 12: 96-102.
- Prasad R. N., Chaturvedi H. C. (1988). Effect of explants on micropropagation of *Chrysanthemum morifolium*. *Biologia Plantarum*, 30: 20-24.
- Prasad R. N., Sharma A. K., Chaturvedi H. C. (1993). Clonal multiplication of *Chrysanthemum morifolium* 'Otome zakura' in long-term culture. *Bangladesh Journal of Botany*, 12: 96-102.
- Rademaker W., de Jong J. (1990). Genetic variation in adventitious shoot formation in *Dendranthema grandiflora* (*Chrysanthemum morifolium*) explants. In: de Jong J. (Ed.). *Integration of in vitro techniques in ornamental plant breeding*, Wageningen: 34-38.
- Ravishankar G. A., Rajasekaran T., Sarma K. S., Venkataraman L.V. (1989). Production of pyrethrins in cultured tissues of pyrethrum *Chrysanthemum cinerariaefolium* Vis. *Pyrethrum Post*, 17: 66-69.
- Renou J. P., Brochard P., Jalouzot R. (1993). Recovery of transgenic chrysanthemum (*Dendranthema grandiflora* Tzvelev) after hygromycin resistance selection. *Plant Science*, 89: 185-197.
- Roberts A. V., Smith E. F. (1990). The preparation *in vitro* of chrysanthemum for transplantation to soil. 1. Protection of roots by cellulose plugs. *Plant Cell, Tissue and Organ Culture*, 21: 129-132.
- Roest S., Bokelmann G. S. (1973). Vegetative propagation of *Chrysanthemum cinerariaefolium in vitro*. *Scientia Horticulturae*, 1: 120-122.
- Roest S., Bokelmann G. S. (1975). Vegetative propagation of *Chrysanthemum morifolium* Ram. *in vitro*. *Scientia Horticulturae*, 3: 317-330.
- Rout G. R., Das P. (1997). Recent trends in the biotechnology of Chrysanthemum: a critical review. *Scientia Horticulturae*, 69: 239-256.

- Rout G. R., Palai S. K., Pandey P., Das P. (1997). Direct plant regeneration of *Chrysanthemum morifolium* Ramat. Deep Pink: influence of explant source, age of explant, culture environment, carbohydrates, nutritional factors and hormone regime. Proceedings of the National Academy of Science India, Section B, Biological Sciences, 67: 57-66.
- Sakai A., Matsumoto T., Hirai D., Niino T. (2000). Newly developed encapsulation-dehydration protocol for plant cryopreservation. Cryo Letters, 21: 53-62.
- Sauvadet M. A., Brochard P., Boccon-Gibco J. (1990). A protoplast-to-plant system in chrysanthemum (*Dendranthema grandiflora* (Ramat.) Kitamura) using stem segments. Plant Cell Reports, 8: 692-695.
- Schwinn K. E., Markham K. R., Given N. K. (1994). Floral flavonoids and their potential for pelargonidin biosynthesis in commercial chrysanthemum cultivars. Phytochemistry, 35: 145-150.
- Schum A., Preil W. (1981). Regeneration of callus from *Chrysanthemum morifolium* mesophyll protoplast. Gartenbauwissenschaft, 46: 91-93.
- Shao H. S., Li J. H., Zheng X. Q., Chen S. C. (1999). Cloning of the *LFY* cDNA from *Arabidopsis thaliana* and its transformation to *Chrysanthemum morifolium*. Acta Botanica Sinica, 41: 268-271.
- Sherman J. M., Moyer J. W., Daub M. E. (1998). A regeneration and *Agrobacterium*-mediated transformation system for genetically diverse chrysanthemum cultivars. Journal of the American Society for Horticultural Science, 123: 189-194.
- Shibli R. A., Smith M. A. L., Spomer L. A. (1992). Osmotic adjustment and growth responses of three *Chrysanthemum morifolium* Ramat. cultivars to osmotic stress induced *in vitro*. Journal of Plant Nutrition, 15: 1373-1381.
- Shinoyama H., Nomura Y., Tsuchiya T., Kazuma T. (1991). Formation and plant regeneration from leaves of chrysanthemum (*Dendranthema grandiflora* Tzvelev.). Journal of the Japanese Society for Horticultural Science, 41 (suppl.): 158.
- Shinoyama H., Nomura Y., Tsuchiya T., Kazuma T. (1997). Direct embryoid formation and plant regeneration from leaves of chrysanthemum (*Dendranthema grandiflora* Tzvelev). Japanese Journal of Breeding, 46: 158.
- Shirasawa N., Iwai T., Nakamura S., Honkura R. (2000). Transformation and transgene expression of chrysanthemum (*Dendranthema grandiflora* (Ramat.) Kitamura). Bulletin of the Miyagi Prefectural Agricultural Research Centre 67: 15-20.
- Staba E. J., Nygaard B. G., Zito S. W. (1984). Light effects on pyrethrum shoot cultures. Plant Cell, Tissue and Organ Culture, 3: 211-214.
- Sun J. X., Li C. P. (1987). Effects of MS medium on root formation off shoot tips from several famed kinds of Chrysanthemum. Acta Scientia Natural University Sunyatseni, 1: 121-122.
- Sutter E., Langhans R. W. (1981). Abnormalities in Chrysanthemum regenerated from long-term cultures. Annals of Botany, 48: 559-568.
- Takatsu Y., Tomotsune H., Kasumi M., Sakuma F. (1998). Differences in adventitious shoot regeneration capacity among Japanese chrysanthemum (*Dendranthema grandiflora* (Ramat.) Kitamura) cultivars and the improved protocol for *Agrobacterium*-mediated genetic transformation. Journal of the Japanese Society for Horticultural Science, 67: 958-964.
- Takatsu Y., Nishizawa Y., Hibi T., Akutsu K. (1999). Transgenic chrysanthemum (*Dendranthema grandiflorum* (Ramat.) Kitamura) expressing a rice chitinase gene shows enhanced resistance to gray mold (*Botrytis cinerea*). Scientia Horticulturae, 82: 113-123.
- Tanaka R., Watanabe K. (1972). Embryological studies in *Chrysanthemum makinoi* and its hybrid crossed with hexaploid. Japanese Journal of Science, 14: 75-84.
- Tanaka K., Kanno Y., Higuchi K., Suzuki M. (1998). Somatic embryo formation and somaclonal variation in chrysanthemum. Journal of the Japanese Society for Horticultural Science, 67: 249.
- Tanaka K., Kanno Y., Kudo S., Suzuki M. (2000). Somatic embryogenesis and plant regeneration in chrysanthemum (*Dendranthema grandiflorum* (Ramat.) Kitamura). Plant Cell Reports, 19: 946-953.
- Tanimoto H., Kagi T. (1990). Culture and regeneration of *Chrysanthemum coronarium* L. leaf protoplasts. Journal of the Japanese Society for Horticultural Science, 59: 252-253.
- Teixeira da Silva J. A. (2002a). Polyamines in the regulation of chrysanthemum and tobacco *in vitro* morphogenic pathways. Propagation of Ornamental Plants, 2: 9-15.
- Teixeira da Silva J. A. (2002b). Antibiotics in successful transformation and in the war against plant tissue culture infection. Newsletter of the International Society for Chemotherapy, 6: 13.
- Teixeira da Silva J. A. (2003a). Thin cell layer technology for induced response and control of rhizogenesis in chrysanthemum. Plant Growth Regulation, 39: 67-76.
- Teixeira da Silva J. A. (2003b). Filter paper significantly affects the morphogenic programmes, and buffers the phytotoxic effect of antibiotics in chrysanthemum and tobacco thin cell layer *in vitro* culture. HortScience,

- 38: (in press).
- Teixeira da Silva J. A. (2003c). Control of chrysanthemum organogenesis by thin cell layer technology. *Asian Journal of Plant Sciences*, 2: 505-514.
- Teixeira da Silva J. A., Fukai S. (2001). The impact of carbenicillin, cefotaxime and vancomycin on chrysanthemum and tobacco TCL morphogenesis and *Agrobacterium* growth. *Journal of Applied Horticulture*, 3: 18-27.
- Teixeira da Silva J. A., Nhut D. T., Tanaka M., Fukai S. (2003). The effect of antibiotics on the *in vitro* growth response of chrysanthemum and tobacco stem transverse thin cell layers (tTCLs). *Scientia Horticulturae*, 97: 397-410.
- Tian X. F., Liu Z. Q., Zhang J. F. (1993). Effect of salt on rooting of *Chrysanthemum* and *Vinca major* shoots *in vitro*. *Acta Horticulturae*, 20: 101-102.
- Tosca A., Delledonne M., Furini A., Belenghi B., Fogher C., Frangi P. (2000). Transformation of Korean chrysanthemum (*Dendranthema zawadskii* x *D. x grandiflorum*) and insertion of the maize autonomous element *Ac* using *Agrobacterium tumefaciens*. *Journal of Genetics and Breeding*, 54: 19-24.
- Tsukamoto Y., Fujime Y. (1971). Flower bud culture of chrysanthemum *in vitro*. *Journal of the Japanese Society for Horticultural Science*, 40: 62-67.
- Urban L. A., Sherman J. M., Moyer J. W., Daub M. E. (1994). High frequency shoot regeneration and *Agrobacterium*-mediated transformation of chrysanthemum (*Dendranthema grandiflora*). *Plant Science*, 98: 69-79.
- van Wordragen M. F., de Jong J., Huitema H. B. M., Dons H. J. M. (1991). Genetic transformation of chrysanthemum using wild type *Agrobacterium* strains; strain and cultivar specificity. *Plant Cell Reports*, 9: 505-508.
- van Wordragen M. F., Ouwerkerk P. B. F., Dons H. J. M. (1992a). *Agrobacterium rhizogenes* mediated induction of apparently untransformed roots and callus in chrysanthemum. *Plant Cell, Tissue and Organ Culture*, 30: 149-157.
- van Wordragen M. F., de Jong J., Schornagel M. J., Dons H. J. M. (1992b). Rapid screening for host-bacterium interactions in *Agrobacterium*-mediated gene transfer to chrysanthemum, by using the GUS-intron gene. *Plant Science*, 81: 207-214.
- Votruba R., Kodytek R. (1988). Investigation of genetic stability in *Chrysanthemum morifolium* 'Blanche Poitevine Suprême' after meristem culture. *Acta Horticulturae*, 226: 311-319.
- Wambugu F. M., Rangan T. S. (1981). *In vitro* clonal multiplication of pyrethrum (*Chrysanthemum cinerariaefolium* Vis.) by micropropagation. *Plant Science Letters*, 22: 219-226.
- Wang S. O., Ma S. S. (1978). Clonal multiplication of Chrysanthemum *in vitro*. *Journal of the Agricultural Association China*, 32: 64-73.
- Watanabe K. (1977). Successful ovary culture and production of F1 hybrids and androgenic haploids in Japanese chrysanthemum species. *Journal of Heredity*, 68: 317-320.
- Widiastoety D. (1987). Preliminary experiment on tissue culture of *Chrysanthemum morifolium*. *Ram. Bulletin of the Penilitian Horticulture*, 15: 231-236.
- Yepes L. C., Mittak V., Pang S. Z., Gonsalves C., Slightom J. L., Gonsalves D. (1995). Biolistic transformation of chrysanthemum with the nucleocapsid gene of tomato spotted wilt virus. *Plant Cell Reports*, 14: 694-698.
- Yepes L. C., Mittak V., Pang S. Z., Gonsalves C., Slightom J. L., Gonsalves D. (1999). *Agrobacterium tumefaciens* versus biolistic-mediated transformation of the chrysanthemum cvs. Polaris and Golden Polaris with nucleocapsid protein genes of three Tosspovirus species. *Acta Horticulturae*, 482: 209-218.
- Yi J., Wang B., Wang X., Duan C., Yang X. (2003). Effect of sound stimulation on roots growth and plasmalemma H⁺-ATPase activity of chrysanthemum (*Gerbera jamesonii*). *Colloids and Surfaces B: Biointerfaces*, 27: 65-69.
- Yiyao L., Bochu W., Xuefeng L., Chuanren D., Sakanishi A. (2002). Effects of sound field on the growth of Chrysanthemum callus. *Colloids and Surfaces B: Biointerfaces*, 24: 321-326.
- Young K. J., Jung P. S., Young U. B., Ho P. C., Soo C. Y., Sheop S. J. (1998). Transformation of chrysanthemum by *Agrobacterium tumefaciens* with three different types of vectors. *Journal of the Korean Society for Horticultural Science*, 39: 360-366.
- Zheng Z. L., Yang Z., Jang J. C., Metzger J. D. (2001). Modification of plant architecture in chrysanthemum by ectopic expression of the tobacco phytochrome *B1* gene. *Journal of the American Society for Horticultural Science*, 126: 19-26.
- Zieg R. G., Zito S. W., Staba E. J. (1983). Selection of high pyrethrin producing tissue cultures. *Planta Medica*, 48: 88-91.
- Zito S. W., Tio C. D. (1990). Constituents of *Chrysanthemum cinerariaefolium* in leaves, regenerated plantlets and callus. *Phytochemistry*, 29: 2533-2534.
- Zito S. W., Zieg R. G., Staba E. J. (1983). Distribution of pyrethrins in oil glands and leaf tissue of *Chrysanthemum cinerariaefolium*. *Planta Medica*, 47: 205-207.