

IDENTIFICATION BY ISOZYME GENE MARKERS OF EUROPEAN PINE SPECIES USED FOR ORNAMENTAL PURPOSES

Apostolos Scaltsoyiannes

Laboratory of Forest Genetics and Plant Breeding, Department of Forestry and Natural Environment,
Aristotle University of Thessaloniki, 54006 Thessaloniki, Greece, Fax: +302310992777,
E-mail: skaltsoy@for.auth.gr

REFERENCES

- Bergmann F., Hattemer H. (1995). Isozyme gene loci and their allelic variation in *Pinus sylvestris* L. and *Pinus cembra* L. *Silvae Genetica*, 44 (5-6): 286-289.
- Bergmann F., Ruetz W. (1987). Short Note: Identifizierung von Hybridlarchensaatgut aus Samenplantagen mit Hilfe eines Isoenzym-Markers. *Silvae Genetica*, 36 (2): 102-105.
- Bletsos F. A., Roupakias D. G., Tsaktsira M. R., Scaltsoyiannes A. B., Thanassouloupoulos C. C. (1998). Interspecific hybrids between three eggplant (*Solanum melongena* L.) cultivars and two wild species (*Solanum torvum* Sw. and *Solanum sisymbriifolium* Lam.). *Plant Breeding*, 117: 159-164.
- Cheliak W. M., Pitel J. A. (1984). Electrophoretic identification of clones in trembling aspen. *Canadian Journal of Forest Research*, 14: 740-743.
- Cheliak W. M., Pitel J. A. (1985). Techniques for starch gel electrophoresis of enzymes from forest tree species. Information report PI-X-42, Petawawa National Forest Institute, 49 pp.
- Conkle M. T., Hodgskiss P. D., Nunnally L. B., Hunter S. C. (1982). Starch gel electrophoresis of conifer seeds: a laboratory manual. In: General Technical Report PSW-64, 18 pp.
- El-Kassaby A. (1981). Genetic interpretation of malate dehydrogenase isozymes in some conifers species. *The Journal of Heredity*, 72: 451-452.
- Huneycutt M., Askew G. R. (1989). Electrophoretic identification of Loblolly Pine-short leaf pine hybrids. *Silvae Genetica*, 38 (3-4): 95-96.
- Krüssmann G. (1983). Manual of cultivated conifers. Timber Press Publishers, 361 pp.
- Little E. L. Jr., Critchfield W. B. (1969). Subdivisions of the genus *Pinus* (Pines). USDA, Forest Service. Miscellaneous Publication No. 1144, 51 pp.
- Mavromatis A., Tsaktsira M., Scaltsoyiannes A., Roupakias D. (in press). Somaclonal variation of *in vitro* cultivated plants of cotton (*Gossypium* sp.p.). Proceedings of the 8th Congress of the Greek Scientific Society for Plant Breeding, Arta 2000: 166-175 (In Greek).
- Mirov N. T. (1967). The genus *Pinus*. The Ronald Press Company, New York, 602 pp.
- Morris R. W., Critchfield W. B., Fowler D. P. (1980). The putative Austrian x Red Pine Hybrid: A test of paternity based on allelic variation at enzyme-specifying loci. *Silvae Genetica*, 29 (3-4): 93-100.
- Moulalis D., Panetsos K. P., Scaltsoyiannes A., Aravanopoulos F. A., Tsaktsira M., Passayiannes G. (1999). Identification and productivity of artificial pine hybrid F₁: *Pinus brutia* x *P. halepensis*. Proceedings of the 8th Congress of the Greek Forest Society, Alexandroupoli 1998: 354-360 (In Greek).
- Nielsen G., Ostergaard H., Johansen H. (1985). Cultivar identification by means of isoenzymes. II. Genetic variation at four enzyme loci in diploid ryegrass. *Zeitschrift für Pflanzenzüchtung*, 94: 74-86.
- Panetsos K. P., Scaltsoyiannes A., Aravanopoulos F. A., Dounavi K., Demetrakopoulos A. (1997). Identification of *Pinus brutia* TEN., *P. halepensis* Mill. and their putative hybrids. *Silvae Genetica*, 46 (5): 253-257.
- Rothe G. M. (1990). Forest Genetic Resources. Information No 18. Food and Agriculture Organization of the United Nations: 1-15.
- Scaltsoyiannes A. (1993). Variation of isozyme patterns of malate dehydrogenase in four subspecies of European black pine (*Pinus nigra* ARNOLD). *Scientific Annals of the Department of Forestry and Natural Environment*, Volume LST, Thessaloniki 1993: 51-61 (In Greek).
- Scaltsoyiannes A. (1999). Allozyme differentiation and phylogeny of Cedar species. *Silvae Genetica*, 48: 61-68.

- Scaltsoyiannes A. (2000). Contribution of malate dehydrogenase on the isozyme analysis and identification of European pine species (*Pinus* sp.). Proceedings of the 8th Congress of the Greek Scientific Society for Plant Breeding, Arta 2000: 166-175 (In Greek).
- Scaltsoyiannes A. (in press). Identification of artificial pine hybrids F_1 : *Pinus brutia* x *P. halepensis* and *Pinus nigra* var. *austriaca* x *P. sylvestris* by means of the gene markers of malate dehydrogenase (MDH). Scientific Annals of the Department of Forestry and Natural Environment, Thessaloniki 2003 (In Greek).
- Scaltsoyiannes A., Moulalis D. (1992). Identification of artificial fir hybrids by biochemical gene markers. *Veliotika*, Volume B, 8-9: 57-60 (In Greek).
- Scaltsoyiannes A., Rohr R., Panetsos K. P., Tsaktsira M. (1994). Allozyme frequency distributions in five European populations of Black pine (*Pinus nigra* Arnold). I. Estimation of genetic variation within and among populations. II. Contribution of isozyme analysis to the taxonomic status of the species. *Silvae Genetica*, 43 (1): 20-30.
- Scaltsoyiannes A., Tsaktsira M., Drouzas A. (1999). Allozyme differentiation in the Mediterranean firs. A first comparative study with phylogenetic implications. *Plant Systematics and Evolution*, 216 (3-4): 289-307.
- Tanksley S. D., Orton T. J. (1983). *Isozymes in plant genetics and breeding*. Elsevier Science Publishers B.V., ISBN 0-444-42228-5, 516 pp.
- Thormann R., Stephan B. R. (1993). Interpretation of isozyme patterns of malate dehydrogenase in Scots pine using two different staining methods. *Silvae Genetica*, 42 (1): 5-8.
- Tsaktsira M., Scaltsoyiannes A. (1998). Isozyme polymorphism of 18 European black pine populations and its contribution to the taxonomic status of the species. Proceedings of the 7th Congress of the Greek Scientific Society for Plant Breeding, Iraklio 1998: 245-254 (In Greek).
- Vidaković M. (1991). *Conifers. Morphology and variation*. Zagreb. 755 pp.