

**IN VITRO PLANT REGENERATION OF ORNAMENTAL CABBAGE**  
**(BRASSICA OLERACEA VAR. LONGATA)**

**Yipeng Zhao<sup>1\*</sup>, Jon Green<sup>2</sup>, Miao Yu<sup>2</sup> and Brian W.W. Grout<sup>2</sup>**

<sup>1</sup>Department of Horticulture, Henan Institute of Science and Technology,  
Xinxiang, Henan 453003, China, \*Fax: + 86-373-304066 \*E-mail: yzhao36@hotmail.com  
<sup>2</sup>Postgraduate School, Writtle College, Lordship Road Chelmsford CM1 3RR, United Kingdom

**REFERENCES**

- Bhalla P. L., Smith N. A. (1998). Comparison of shoot regeneration potential from seedling explants of Australian cauliflower (*Brassica oleracea* var. *botrytis*) varieties. *Australian Journal of Agricultural Research*, 49 (8): 1261-1266.
- Grout B. W. W. (1990). Meristem tip culture. *In: Pollard J. W., Walker J. M. (Eds.). Plant Cell and Tissue Culture. The Humana Press Inc, Clifton, New Jersey: 81-91.*
- Grout B. W. W., Crisp P. (1977). Practical aspects of the propagation of cauliflower by meristem culture. *Acta Horticulturae*, 78: 289-296.
- Grout B. W. W., Crisp P. (1980). The origin and nature of shoots propagated from cauliflower roots. *Journal of Horticultural Science*, 55 (1): 65-70.
- Grout B. W. W., Westcott R. J., Henshaw G. G. (1977). Scanning electron microscope studies of multiple shoot production by meristem tip cultures of *Solanum x curtilobum*. *Annals of Botany*, 41: 1113-1116.
- Hwang S. J. (2004). An efficient *in vitro* protocol for propagation of *Zanthoxylum piperitum* DC. *Propagation of Ornamental Plants*, 2: 37-40.
- Jain S. M. (2001). Tissue culture-derived variation in crop improvement. *Euphytica*, 118: 153-166.
- Loiseau J., Michaux-Ferrere N., Deunff Y. L. (1998). Histological investigations of somatic embryogenesis in peas. *Plant Physiology and Biochemistry*, 36: 683-687.
- Murashige T., Skoog F. (1962). A revised medium for rapid rowth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- Msikita W., Skirvin R. M. (1989). *In vitro* regeneration from hypocotyls and seedling cotyledons of tronchuda (*Brassica oleracea* var. *tronchuda* Bailey). *Plant Cell, Tissue and Organ Culture*, 19: 159-165.
- Narasimhulu S. B., Chopra V. L. (1988). Species-specific shoot regeneration response of cotyledonary explants of Brassicas. *Plant Cell Reports*, 7: 473-497.
- Pua E. C., Teo S. H., Loh C. S. (1996). Interactive role of ethylene and polyamines on shoot regenerability of Chinese cabbage (*Brassica alboglabra* Bailey) *in vitro*. *Journal of Plant Physiology*, 149: 138-148.
- Pua E. C., Trinh T. H., Chua N. H. (1989). High frequency plant regeneration from stem explants of *Brassica alboglabra* Bailey *in vitro*. *Plant Cell, Tissue and Organ Culture*, 17: 143-152.
- Sretenović-Rajčić T., Mijatović M., Vinterhalter D. (2002). *In vitro* culture as a tool for improvement of cabbage cultivars in Yougoslavia. *Acta Horticulturae*, 579: 209-213.