

**PROPAGATION OF JAPANESE RED PINE (*PINUS DENSIFLORA* ZIEB. ET ZUCC.)
VIA SOMATIC EMBRYOGENESIS**

Emilio Maruyama*, Yoshihisa Hosoi, and Katsuaki Ishii

Department of Molecular and Cell Biology, Forestry and Forest Products Research Institute, P.O. Box 16,
Tsukuba Norinckenkyudanchi-Nai, Ibaraki, 305-8687 Japan

*Fax: +81-29-873-1542, *E-mail: tsumaruy@ffpri.affrc.go.jp

REFERENCES

- Becwar M. R., Nagmani R., Wann S. R. (1990). Initiation of embryogenic cultures and somatic embryo development in loblolly pine (*Pinus taeda*). *Canadian Journal of Forest Research*, 20: 810-817.
- Duncan D. B. (1995). Multiple range and multiple F test. *Biometrics*, 11: 1-42.
- Garin A., Isabel N., Plourde A. (1998). Screening of large numbers of seed families of *Pinus strobus* L. for somatic embryogenesis from immature and mature zygotic embryos. *Plant Cell Reports*, 18: 37-43.
- Hosoi Y., Ishii K. (2001). Somatic embryogenesis and plantlet regeneration in *Pinus armandii* var. *amamiana*. In: Morohoshi N., Komamine A. (Eds.). *Molecular breeding of woody plants*, Elsevier Science: 313-318.
- Ishii K., Maruyama E., Hosoi Y. (2001). Somatic embryogenesis of Japanese conifers. In: Morohoshi N., Komamine A. (Eds.). *Molecular breeding of woody plants*, Elsevier Science: 297-304.
- Jain S. M., Gupta P. K., Newton R. J. (1995). *Somatic embryogenesis in woody plants*, Vol. 3, Gymnosperms. Kluwer Academic Publishers, 388 pp.
- Jain S. M., Gupta P. K. (2005). *Protocol for somatic embryogenesis in woody plants*. Springer, 585 pp.
- Klimaszewska K., Smith D. R. (1997). Maturation of somatic embryos of *Pinus strobus* is promoted by a high concentration of gellan gum. *Physiologia Plantarum*, 100: 949-957.
- Lelu M. A., Bastien C., Drugeault A., Gouez M. L., Klimaszewska K. (1999). Somatic embryogenesis and plantlet development in *Pinus sylvestris* and *Pinus pinaster* on medium with and without growth regulators. *Physiologia Plantarum*, 105: 719-728.
- Li X. Y., Huang F. H., Gbur E. E. (1997). Polyethylene glycol-promoted development of somatic embryo in loblolly pine (*Pinus taeda* L.). *In Vitro Cellular and Development Biology-Plant*, 33: 184-189.
- Li X. Y., Huang F. H., Murphy J. B., Gbur E. E. (1998). Polyethylene glycol and maltose enhance somatic embryo maturation in loblolly pine (*Pinus taeda* L.). *In Vitro Cellular and Developmental Biology-Plant*, 34: 22-26.
- Maruyama E., Tanaka T., Hosoi Y., Ishii K., Morohoshi N. (2000). Embryogenic cell culture, protoplast regeneration, cryopreservation, biolistic gene transfer and plant regeneration in Japanese cedar (*Cryptomeria japonica* D. Don). *Plant Biotechnology*, 17: 281-296.
- Maruyama E., Hosoi Y., Ishii K. (2002). Somatic embryogenesis in Sawara cypress (*Chamaecyparis pisifera* Sieb. et Zucc.) for stable and efficient plant regeneration, propagation and protoplast culture. *Journal of Forest Research*, 7: 23-34.
- Maruyama E., Ishii K., Hosoi Y. (2005). Efficient plant regeneration of Hinoki cypress (*Chamaecyparis obtusa* Sieb. et Zucc.) via somatic embryogenesis. *Journal of Forest Research*, 10: 73-77.
- Smith D. R. (1996). Growth medium. United States Patent 5 565 355.
- Taniguchi T. (2001). Plant regeneration from somatic embryos in *Pinus thunbergii* (Japanese black pine) and *Pinus densiflora* (Japanese red pine). In: Morohoshi N., Komamine A. (Eds.). *Molecular breeding of woody plants*, Elsevier Science: 318-324.
- Walter C. (2004). Genetic engineering in conifer forestry: technical and social considerations. *In Vitro Cellular and Developmental Biology-Plant*, 40: 434-441.