

**IN VIVO DIFFERENTIATION OF ADVENTITIOUS SHOOTS IN DECAPITATED PLANTS  
OF POINSETTIA (*EUPHORBIA PULCHERRIMA* WILLD EX KLOTSCH)**

**Malene D. Nielsen<sup>1</sup>, Priyavadan A. Joshi<sup>2</sup>, and Arne Skytt Andersen<sup>1\*</sup>**

<sup>1</sup>Department of Agricultural Sciences, <sup>2</sup>Department of Plant Biology, The Royal Veterinary and Agricultural University, Thorvaldsensvej 40, DK-1871 Frederiksberg C, Denmark,

\*Fax: +45 3528 3400, \*E-mail: asa@kvl.dk

**REFERENCES**

- Asseyeva T. (1927). Bud mutations in the potato and their chimerical nature. *Journal of Genetics*, 19: 1-26.
- Bateson W. (1921). Root cuttings and chimaeras 2. *Journal of Genetics*, 11: 91-97.
- Bergann F., Bergann L. (1982). Zur Entwicklungsgeschichte des Angiospermenblattes 1. Über periklinalchimären bei *Peperomia* und ihre experimentelle Entmischung und Umlagerung. *Biologisches Zentralblatt* 101: 485-502.
- Broertjes C., Keen A. (1980). Adventitious shoots: do they develop from one cell? *Euphytica*, 29: 73-87.
- Broertjes C., Leffring L. (1972). Mutation breeding of kalanchoë. *Euphytica*, 21: 415-423.
- Broertjes C., Lock C. A. M. (1985). Radiation-induced low-temperature tolerant solid mutants of *Chrysanthemum morifolium* Ram. *Euphytica*, 34: 97-103.
- Broertjes C., van Harten A. M. (1985a.). General Introduction. *In*: Broertjes, C., van Harten A. M. (Eds.). *Developments in Crop Science 12: Applied Mutation Breeding for Vegetatively Propagated Crops*. Elsevier: 3-13.
- Broertjes C., van Harten A. M. (1985b). Single cell origin of adventitious buds. *Euphytica*, 34: 93-95.
- Broertjes C., van Harten A. M. (1988). Shoot apices: Organization and post-irradiation behavior. *In*: Broertjes C., van Harten A. M. (Eds.). *Developments in crop science 12: Applied mutation breeding for vegetatively propagated crops*. Elsevier: 29-44.
- Christianson M. L., Warnick D. A. (1988). Organogenesis *in vitro* as a developmental process. *HortScience*, 23: 515-519.
- Fink S. (1999). Pathological and regenerative plant anatomy. *Encyclopedia of plant anatomy*. Volume XIV/6, Bornträger, Berlin: 371-404.
- Graham E. (1997). Staining Tomato fruit cuticle and exocarp tissues. *Biotechnology and Histochemistry*, 72: 119-122.
- Graham E., Joshi P. A. (1995). Novel fixation of plant tissue, staining through paraffin with alcian blue and hematoxylin, and improved slide preparation. *Biotechnology and Histochemistry*, 70: 263-266.
- Graham E. T., Trentham W. R. (1998). Staining paraffin extracted, alcohol rinsed and air dried plant tissue with an aqueous mixture of three dyes. *Biotechnology and Histochemistry*, 73: 178-185.
- Lee I. M., Klopmeier M., Bartoszyk I. M., Gundersen-Rindal D. E., Chou T-S., Thomson K. L., Eisenreich R. (1997). Phytoplasma induced free-branching in commercial poinsettia cultivars. *Nature Biotechnology*, 15: 178-182.
- Lo K. H., Giles K. L., Sawhney V. K. (1997). Histological changes associated with acquisition of competence for shoot regeneration in leaf discs of *Saintpaulia ionantha* x *confusa* hybrids (African violet) cultured *in vitro*. *Plant Cell Reports*, 16: 421-425.
- Maede E., Thorpe T. A. (1979). Shoot histogenesis in tobacco callus cultures. *In vitro*, 15: 415-424.
- Marcotrigiano M. (1986). Origin of adventitious shoots regenerated from cultured tobacco leaf tissue. *American Journal of Botany*, 73: 1541-1547.
- Neilson-Jones W. (1969). *Plant chimeras*. Methuen and Co., 123 pp.
- Nielsen M. D., Farestveit B., Andersen A. S. (2003). Adventitious shoot development from decapitated plants of periclinal chimeric poinsettia plants (*Euphorbia pulcherrima* Wild ex Klotsch). *Europaen Journal of Horticultural Science*, 68: 161-168.
- Novak F. J. (1991). *In vitro* mutation system for crop improvement. *In*: *Plant Mutation Breeding for Crop Improvement*. Volume 2. International Atomic Energy Agency: 327-342.

- Stewart R. N. (1960). Inheritance of bract color in poinsettia. *Journal of Heredity*, 51: 175-177.
- Stewart R. N. (1965). The origin and transmission of a series of plastogene mutants in *Dianthus* and *Euphorbia*. *Genetics*, 52: 925-947.
- Stewart R. N., Arisumi A. (1966). Genetic and histogenic determination of pink bract color in poinsettia. *Journal of Heredity*, 57: 216-220.
- Stewart R. N., Dermen H. (1970). Determination of number and mitotic activity of shoot apical initial cells by analysis of mericlinal chimeras. *American Journal of Botany*, 57: 816-826.
- Szymkowiak E. J., Sussex I. M. (1992). The internal meristem layer (L3) determines floral meristem size and carpel number in tomato periclinal chimeras. *Plant Cell*, 4: 1089-1100.
- Tian H-C., Marcotrigiano M. (1993). Origin and development of adventitious shoot meristems initiated on plant chimeras. *Developmental Biology*, 155: 259-269.
- Tilney-Bassett R. A. E. (1986). *Plant Chimeras*. E. Arnold Publishers, London, 199 pp.