

AUTONOMOUS SHOOT PRODUCTION IN PAWPAW (*ASIMINA TRILOBA* (L.) DUNAL) ON PLANT GROWTH REGULATOR FREE MEDIA

Robert Geneve^{1*}, Sharon Kester¹, and Kirk Pomper²

¹Department of Horticulture University of Kentucky, Lexington, KY 40546, USA,

*Fax: + 1 859-257-2859, *E-mail: RGeneve@uky.edu

²Kentucky State University, Frankfort, KY 40601, USA

REFERENCES

- Brand M. H., Kiyomoto R. (1997). Induction of tissue proliferation-like characteristics in *in vitro* cultures of *Rhododendron* 'Montego'. *HortScience*, 32: 989-994.
- Catterou M., Debois F., Smets R., Vanriet S., Kichey T., Onckelen H. V., Sangwan-Norreel B. S., Sangwan R. S. (2002). hoc: an *Arabidopsis* mutant overproducing cytokinins and expressing high *in vitro* organogenic capacity. *Plant Journal*, 30: 273-287.
- Datta M. M., Majumder A., Jha S. (2006). Organogenesis and plant regeneration in *Taxus wallichiana* (Zucc.). *Plant Cell Reports*, 25: 11-18.
- Finneseth C. H., Layne D. R., Geneve R. L. (1998). Requirements for seed germination in North American pawpaw [*Asimina triloba* (L.) Dunal]. *Seed Science and Technology*, 26: 471-480.
- Finneseth C. L. H., Layne D. R., Geneve R. L. (2000). Establishment of North American pawpaw [*Asimina triloba* (L.) Dunal] shoots *in vitro* from mature and juvenile explants. *Acta Horticulturae*, 520: 97-102.
- Gautheret R. J. (1942). Hetero-auxines et cultures de tissus vegetaux. *Bulletin de la Société Chimique*, 24: 13-46.
- Geneve R. L., Pomper K. W., Kester S. T., Egilla J. N., Finneseth C. L. H., Crabtree S., Layne D. R. (2003). Propagation of Pawpaw - a Review. *HortTechnology*, 13: 428-433.
- Hawes M. C., Sharpe D. Z., Plata M. I., Pueppke S. G., Chourey P. S. (1985). Auxin-independent growth of maize tissue culture cells. *Plant Science*, 40: 197-202.
- Henderson J. H. M. (1954). The changing nutritional pattern from normal to habituated sunflower callus tissue *in vitro*. *Annee Biologique*, 30: 329-348.
- Hewelt A., Prinsen E., Thomas M., Van Onckelen H., Meins F. Jr. (2000). Ectopic expression of maize *knotted1* results in the cytokinin-autotrophic growth of cultured tobacco tissues. *Planta*, 210: 884-889.
- Huetteman C. A., Preece J. E. (1993). Thidazuron: a potent cytokinin for woody plant tissue culture. *Plant Cell, Tissue and Organ Culture*, 33: 105-119.
- Hwang I., Sheen J. (2001). Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature*, 413: 383-389.
- Kohumura H., Sakai A., Chokyu S., Yakuwa T. (1992). Cryopreservation of *in vitro*-cultured bud clusters of asparagus (*Asparagus officinalis* L. cv. Hiroshimagreen (2n = 30). *Plant Cell Reports*, 11: 433-437.
- Layne D. R. (1996). The pawpaw [*Asimina triloba* (L.) Dunal]: A new fruit crop for Kentucky and the United States. *HortScience*, 31: 777-784.
- Medford J. I., Horgan R., El-Sawi Z., Klee H. J. (1989). Alterations of endogenous cytokinins in transgenic plants using a chimeric isopentenyl transferase gene. *Plant Cell*, 1: 403-413.
- Maene L., Debergh P. (1985). Liquid medium additions to establish tissue cultures to improve elongation and rooting *in vivo*. *Plant Cell, Tissue and Organ Culture*, 5: 23-33.
- Meins F. Jr. (1989). Habituation: heritable variation in the requirement of cultured plant cells for hormones. *Annual Review of Genetics*, 23: 395-408.
- Meins F. Jr., Binns A. (1977). Epigenetic variation of cultured somatic cells. Evidence for gradual changes in requirements for factors promoting cell division. *Proceedings of the National Academy of Sciences of USA*, 74: 2928-2932.
- Meins F. Jr., Thomas M. (2003). Meiotic transmission of epigenetic changes in the cell-division factor requirements of plant cells. *Development*, 130: 6201-6208.

- Murashige T., Skoog F. A. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- Osakabe Y., Yiyata S., Urae T., Seki M., Shinozaki K., Yamaguchi-Shinozaki K. (2002). Overexpression of *Arabidopsis* response regulators, ARR4/ATRR1/IBC7 and ARR8/ATRR3, alters cytokinin responses differentially in the shoot and in callus formation. *Genes and Development*, 16: 479-489.
- Pischke M. S., Huttlin E. L., Hegeman A. D., Sussman M. R. (2006). A transcriptome-based characterization of habituation in plant tissue culture. *Plant Physiology*, 140: 1255-1278.
- Quoirin M., da Silva M. C., Martins K. G., de Oliveira D. E. (2001). Multiplication of juvenile black wattle by microcuttings. *Plant Cell, Tissue and Organ Culture*, 66: 199-205.
- Rodríguez R., Diaz-Sala C., Cuozzo L., Ancora G. (1991). Pear *in vitro* propagation using a double-phase culture system. *HortScience*, 26: 62-64.
- Sujatha M., Reedy T. P. (1998). Differential cytokinin effects on the stimulation of in vitro shoot proliferation from meristematic explants of castor (*Ricinus communis* L.). *Plant Cell Reports*, 17: 561-566.
- Weatherland M. A., Burdon J., Henshaq G. G. (1978). Some effects of activated charcoal as an additive to plant tissue culture media. *Zeitschrift für Pflanzenphysiologie*, 89: 141-144.