

IMPROVING ROOTING UNIFORMITY IN ROSE CUTTINGS

Hendrik-Jan van Telgen*, Barbara Eveleens, and Nieves García

Wageningen UR Glasshouse Horticulture, 2665 MV Bleiswijk, The Netherlands,

*Fax: + 31 174 636 835, *E-mail: hendrikjan.vantelgen@wur.nl

REFERENCES

- Bertram L. (1992). The physiological background for regulation of adventitious root formation by the irradiance to stock plants: a hypothesis. *Acta Horticulturae*, 314: 291-300.
- Bradford K. J., Trewavas A. J. (1994). Sensitivity thresholds and variable time scales in plant hormone action. *Plant Physiology*, 105: 1029-1036.
- Bredmose N., Hansen J. (1995). Regeneration, growth and flowering of cut rose cultivars as affected by propagation material and method. *Scientia Horticulturae*, 64: 103-111.
- Bredmose N., Hansen J. (1996). Topophysis affects the potential of axillary bud growth, fresh biomass accumulation and specific fresh weight in single-stem roses (*Rosa hybrida* L.). *Annals of Botany*, 78: 215-222.
- Bredmose N. (1997). Chronology of three physiological development phases of single-stemmed rose (*Rosa hybrida* L.) plants in response to increment in light quantum integral. *Scientia Horticulturae*, 69: 107-115.
- Costa J. M., Challa H., Van Meeteren U., van de Pol P. A. (2001). Photosynthates: mainly stored and yet limiting in propagation of rose cuttings. *Acta Horticulturae*, 547: 167-174.
- Davies P. (1995). The plant hormone concept: concentration, sensitivity and transport. In: Davies P. (Ed.). *Plant Hormones: Physiology, Biochemistry and Molecular Biology*, 2nd Edition. Kluwer Academic Publishers: 13-38.
- De Klerk G. J., Van der Krieken W., De Jong J. C. (1999). The formation of adventitious roots: new concepts, new possibilities. *In vitro Cellular and Developmental Biology-Plant*, 35: 189-199.
- De Kreijl C., Voogt W., van den Bos A. L., Baas R. (1997). Voedingsoplossingen voor de teelt van roos in gesloten teeltsystemen. Brochure 4. Proefstation voor Bloemisterij en Glasgroenten, 35 pp (in Dutch).
- De Vries D. P., Dubois L. A. M. (1992). Variation in the shoot production of 'Sonia' cut rose plants, originating from grafting different nodal positions. *Gartenbauwissenschaft*, 57: 64-68.
- Dubois L. A. M., de Vries D. O. (1985). Aanwijzingen voor opkweek goed kasrozenstek. *Vakblad voor de Bloemisterij*, 45: 26-31 (in Dutch).
- Fought D. L., Anderson R. G. (1989). Greenhouse production of cut roses from single node cuttings. *Hort Science*, 23: 322.
- Garcia Victoria N., Eveleens B., Van Telgen H. J. (2007a). Development of a phase dependent growth strategy for mobile rose cultivation systems. *Acta Horticulturae*, 175: 137-148.
- Garcia Victoria N., Eveleens B., Van Telgen H. J., van Weel P. (2007b). Critical success factors for robotized rose harvesting. *Acta Horticulturae*, 175: 33-42.
- Kenney G., Sudi J., Blackman G. E. (1969). The uptake of growth substances. XIII. Differential uptake of indol-3yl-acetic acid through the epidermal and cut surfaces of etiolated stem segments. *Journal of Experimental Botany*, 20: 820-840.
- Moe R. (1973). Propagation, growth and flowering of potted roses. *Acta Horticulturae*, 31: 35-49.
- Telgen H. J. van, van Mil A., Kunneman B. (1992a). Effect of propagation and rooting conditions on acclimatization of micropropagated plants. *Acta botanica neerlandica*, 41 (4): 453-459.
- Telgen H. J. van, Elagöz V., van Mil A., Paffen A., de Klerk G. J. (1992b). Role of plant hormones in lateral bud growth of rose and apple *in vitro*. *Acta Horticulturae*, 319: 137-142.
- Wang Q., Skytt Andersen A. (1989). Propagation of *Hibiscus rosa-sinensis*: relations between stock plant cultivar, age, environment and growth regulator treatments. *Acta Horticulturae*, 251: 289-309.