

**MORPHOLOGICAL FEATURES OF PROEMBRYOGENIC MASSES IN RELATION TO THE  
ABILITY OF SOMATIC EMBRYO FORMATION IN CONIFERS**

**Remi Nakagawa<sup>1,3\*</sup>, Shinjiro Ogita<sup>2</sup>, Ugai Watanabe<sup>3</sup>, Takafumi Kubo<sup>1</sup>, and Ryo Funada<sup>1</sup>**

<sup>1</sup>Faculty of Agriculture, Tokyo University of Agriculture and Technology, Fuchu, Tokyo, Japan,

\*Fax: + 81-42-334-5700, \*E-mail: funada@cc.tuat.ac.jp

<sup>2</sup>Biotechnology Research Center, Toyama Prefectural University, Imizu, Toyama, Japan

<sup>3</sup>High technology Research Center, Chiba Institute of Technology, Narashino, Chiba, Japan

**REFERENCES**

- Aitken-Christie J., Thorpe T. A. (1984). Clonal propagation: Gymnosperms. *In*: Vasil I. K. (Ed.). Cell culture and somatic cell genetics of Plants. Vol. 1, Academic Press Inc, London, New York and San Francisco: 82-95.
- Campbell R. A., Durzan D. J. (1975). Induction of multiple buds and needles in tissue cultures of *Picea glauca*. *Canadian Journal of Botany*, 53: 1652-1657.
- Filonova L. H., Bozhkov P. V., von Arnold S. (2000). Developmental pathways of somatic embryogenesis in *Picea abies* as revealed by time-lapse tracking. *Journal of Experimental Botany*, 51: 249-264.
- Nagl W. (1990). Translocation of putrescine in the ovule suspensor and embryo of *Phaseolus coccineus*. *Journal of Plant Physiology*, 136: 587-591.
- Nakagawa R., Ogita S., Kubo T., Funada R. (2006). Effect of polyamines and L-ornithine in the development of proembryogenic masses of *Cryptomeria japonica*. *Plant Cell, Tissue and Organ Culture*, 85: 229-234.
- Stasolla C., Yeung E. C. (2003). Recent advances in conifer somatic embryogenesis: improving somatic embryo quality. *Plant Cell, Tissue and Organ Culture*, 74: 15-35.
- Yeung E. C. (1980). Embryogeny of *Phaseolus*: the role of suspensor. *Zeitschrift für Pflanzenphysiologie*, 96: 17-28.