

**DIRECT AND INDIRECT *IN VITRO* ORGANOGENSIS OF
LEUCOCORYNE PURPUREA (ALLIACEAE) A CHILEAN ORNAMENTAL GEOPHYTE**

Luis H. Escobar^{1*}, Miguel Jordan², Eduardo Olate¹, Luis Barrales³, and Marlene Gebauer¹

¹Department of Plant Science, Pontificia Universidad Católica de Chile, Santiago de Chile, Chile,
*Fax: + 56-2 552-0780, *E-mail: eolate@uc.cl

²Department of Ecology, Pontificia Universidad Católica de Chile, Santiago de Chile, Chile

³Biometry Unit, Pontificia Universidad Católica de Chile, Santiago de Chile, Chile

REFERENCES

- Arriagada L., Mansur L., Zöllner O., Verdugo G., De la Cuadra C., Chellet V., Vergara R., Quiroz M. (2000). Creación *ex-situ* de una colección del género *Leucocoryne*. *Gayana Botánica*: 57: 90 (in Spanish)
- Ayabe M., Sumi S. (1998). Establishment of a novel tissue culture method, stem-disc culture, and its practical application to micropropagation of garlic (*Allium sativum* L.). *Plant Cell Reports*, 17: 773-779.
- Barringer S. A., Mohamed-Yasseen Y., Schloupt R. M., Splittstoesser W. E. (1996). Regeneration of *Allium* spp. *in-vitro* by slicing the basal plate. *Journal of Vegetable Crop Production*, 2: 27-33.
- Bohanec B., Jakse M. (1999). Variations in gynogenic response among long-day onion (*Allium cepa* L.) accessions. *Plant Cell Reports*, 18: 737-742.
- De Hertogh A. A., Le Nard M. (1993). Botanical aspects of flower bulbs. *In: De Hertogh A. A., Le Nard M. (Eds.). The physiology of flower bulbs*. Elsevier, Amsterdam, Holland: 7-20.
- Haque M. S., Wada T., Hattori K. (1997). High frequency shoot regeneration and plantlet formation from root tip from garlic. *Plant Cell, Tissue and Organ Culture*, 50: 83-89.
- Haque M. S., Wada T., Hattori K. (1999). Anatomical changes during *in vitro* direct formation of shoot bud from root tips in garlic (*Allium sativum* L.). *Plant Production Science*, 2: 146-153.
- Kim E. K., Hahn E. J., Murthy H. N., Paek K. Y. (2003). High frequency of shoot multiplication and bulblet formation of garlic in liquid cultures. *Plant Cell, Tissue and Organ Culture*, 73: 231-236.
- Luciani G. F., Marinangeli P. A., Curvetto N. R. (2001). Increasing nitrate/ammonium ratio for improvement of garlic micropropagation. *Scientia Horticulturae*, 87: 11-20.
- Mansur L., Zöllner O., Riedemann P. G., Harrison C. (2002). *Leucocoryne*, un género nativo chileno y su uso como planta de jardín. Manual № 1. Facultad de Agronomía. Universidad Católica de Valparaíso. Valparaíso, Chile, 49 pp. (in Spanish).
- Mohamed-Yasseen Y. (2002). *In-vitro* callus formation and plant regeneration from several onion (*Allium cepa* L.) explants. *Bulletin of the Faculty of Agriculture, University of Cairo*, 53: 289-298.
- Mohamed-Yasseen Y., Nasr M. I. (2003). Callus induction and plant regeneration from Kurrat (*Allium ampeloprasum* var. *kurrat*) and Leek (*Allium ampeloprasum* var. *porrum*). *Bulletin of the Faculty of Agriculture, University of Cairo*, 54: 495-508.
- Murashige T., Skoog F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- Pevalek-Kozlina B., Prolic M. (2001). Differentiation and morphogenetic potencial of *Allium commutatum* Guss. callus tissue. *Acta Botanica Croatica*, 60 (2): 169-175.
- SAS/STAT User's Guide. (1999). SAS Institute Inc., Cary, NC. USA.
- Sata S. J., Bagataria S. B., Thaker V. S. (2001). Induction of direct somatic embryogenesis in garlic (*Allium sativum*). *Methods in Cell Science*, 22: 299-304.
- Soon-Seob K., De-Ping-Guo Jung D. C., Soon-Tae K. (2003). Multiple shoot regeneration and *in-vitro* bulbet formation from garlic callus. *Journal of Plant Biotechnology*, 5: 95-99.
- Squeo F. A., Arancio G., Gutiérrez J. R. (2001). Libro rojo de la flora nativa y de los sitios prioritarios para su conservación: Región de Coquimbo. Ediciones de la Universidad de la Serena, Santiago, 372 pp. (in Spanish).
- Susek A., Javornik B., Bohanec B. (2002). Factors affecting direct organogenesis from flower explant of *Allium giganteum*. *Plant Cell, Tissue and Organ Culture*, 68: 27-33.
- Toaima N., Novak E., Schumann G. (2003). Callus induction from different explants of commercial cultivars of Leek, *Allium ampeloprasum* var. *porrum* L. *Acta Horticulturae*, 597: 303-309.
- Zar J. C. (1984). *Biostatistical Analysis*. 2nd Edition. Prentice Hall. Englewood Cliffs, NJ, USA. 718 pp.
- Zheng S. J., Henken B., Krens F. A., Keizer P. (2003). The development of an efficient cultivar-independent plant regeneration system from callus derived from both apical and non-apical root segments of garlic (*Allium sativum* L.). *In Vitro Cellular and Developmental Biology-Plant*, 39: 288-292.
- Zheng S. J., Henken B., Sofiari E., Keizer P., Kik C., Krens F. A. (1999). Effect of cytokinins and lines on plant regeneration from long-term callus and suspension cultures of *Allium cepa* L. *Euphytica*, 108: 83-90.