

**THE ROLE AND SIGNIFICANCE OF POLYPHENOLS IN ADVENTITIOUS ROOTING IN *PRUNUS*
REGARDING PHYSIOLOGICAL STATUS OF THE STOCK PLANTS**

Gregor Osterc*, Mateja Štefančič, and Franci Štampar

University of Ljubljana, Biotechnical Faculty, Department of Agronomy, Institute for Fruit Growing,
Viticulture and Vegetable Growing, 101 Jamnikarjeva str., 1000 Ljubljana, Slovenia,
Fax: + 386 1 423 10 88, *E-mail: gregor.osterc@bf.uni-lj.si

REFERENCES

- Faivre-Rampant O., Charpentier J.-P., Kevers C., Dommès J., van Onckelen H., Jay-Allemand C., Gaspar T. (2002). Cuttings of the non-rooting *rac* tobacco mutant overaccumulate phenolic compounds. *Functional Plant Biology*, 29: 63-71.
- Fernandez-Lorenzo J. L., Rigueiro A., Ballester A. (1999). Polyphenols as potential markers to differentiate juvenile and mature chestnut shoot cultures. *Tree Physiology*, 19: 461-466.
- Haffner V., Enjalric L., Lardet L., Carron M. P. (1991). Maturation of woody plants: a review of metabolic and genomic aspects. *Annales des Sciences Forestières*, 48: 615-630.
- Hartmann H. T., Kester D. E., Davies F. T. Jr., Geneve R. L. (1997). *Plant propagation-Principles and Practices*. 6th ed., Prentice Hall, New Jersey, USA, 770 pp.
- Leakey R. R. B. (2004). Physiology of vegetative reproduction. *In: Burley J., Evans J., Youngquist J. A. (Eds.). Encyclopaedia of Forest Sciences*, Elsevier, Academic Press: 1655-1668.
- Nag S., Saha K., Choudhuri M. A. (2001). Role of auxin and polyamines in adventitious root formation in relation to changes in compounds involved in rooting. *Journal of Plant Growth Regulation*, 20: 182-194.
- Osterc G., Štampar F. (2003). Propagation of the ornamental cherry *Prunus subhirtella* var. *autumnalis* by soft-wood cuttings. *Propagation of Ornamental Plants*, 3: 47-49.
- Osterc G., Štefančič M., Solar A., Štampar F. (2007). Potential involvement of flavonoids in the rooting response of chestnut hybrid (*Castanea crenata* × *Castanea sativa*) clones. *Australian Journal of Experimental Agriculture*, 47: 96-102.
- Osterc G., Štefančič M., Solar A., Štampar F. (2008). Phenolic acids affect rooting in chestnut hybrid clones (*Castanea crenata* × *Castanea sativa*). *Acta Agriculturae Scandinavica. Section B: Soil & Plant Science*, 58: 162-168.
- Scalbert A., Monties B., Favre J.-M. (1988). Polyphenols of *Quercus robur*: adult tree and *in vitro* grown calli and shoots. *Phytochemistry*, 27: 3483-3488.
- Spethmann W. (1997). Autovegetative Gehölzvermehrung. *In: Krüssmann G. (Ed.). Die Baumschule*, Parey Verlag: 382-449 (in German).
- Tarragó J., Sansberro P., Filip R., López P., Gonzáles A., Luna C., Mroginski L. (2005). Effect of leaf retention and flavonoids on rooting of *Ilex paraguariensis* cuttings. *Scientia Horticulturae*, 103: 479-488.
- Trobec M., Štampar F., Veberič R., Osterc G. (2005). Fluctuations of different endogenous phenolic compounds and cinnamic acid in the first days of the rooting process of cherry rootstock 'Gisela 5' leafy cuttings. *Journal of Plant Physiology*, 162: 589-597.