

## THE EFFECTS OF CARBON DIOXIDE ON ROOTING OF WOODY PLANTS

**Katsuaki Ishii\*, Yoshihisa Hosoi, and Emilio Maruyama**

Forestry and Forest Products Research Institute, Department of Molecular and Cell Biology, Matsunosato 1,  
Tsukuba, Ibaraki 305-8687, Japan,

\*Fax: +81-298-873-1542, \*E-mail: katsuaki@ffpri.affrc.go.jp

### REFERENCES

- Fournioux J. C., Bessis R. (2004). Use of carbon dioxide enrichment to obtain adult morphology of grapevine *in vitro*. *Plant Cell Tissue and Organ Culture*, 33: 51-57.
- Idso S. B. (1999). The long-term response of trees to atmospheric CO<sub>2</sub> enrichment. *Global Change Biology* 5:493-495.
- Ishii K. (1993). Screening of tissue culture conditions of Hinoki cypress and Japanese black pine. *Bulletin of the Forestry and Forest Products Research Institute*, 365: 131-167.
- Kawaoka A., Tanabe T., Shimizu K., Fujii Y., Hayashi K., Schiller C., Murakami K. (2007). Clonal plantation of *Eucalyptus globules* in Western Australia. Abstract of IUFRO Tree Biotechnology 2007, SIX.3.
- Nagae S., Takamura T., Tanabe T., Murakami A., Murakami K., Tanaka M. (1996). *In vitro* shoot development of *Eucalyptus citriodora* on rockwool in the film culture vessel under CO<sub>2</sub> enrichment. *Journal of Forest Research* 1: 227-230.
- Nowak J. (2004). The effects of rooting media, CO<sub>2</sub> enrichment, P-nutrition and mycorrhizal inoculation on rooting and growth of *Osteospermum*. *Acta Horticulturae*, 644: 589-593.
- White P. R. (1943). Nutrient deficiency studies and an improved inorganic nutrient for cultivation of excised tomato roots. *Growth*, 7:53-65.
- Yahara T., Ohba H., Murata J., Iwatsuki K. (1987). Taxonomic review of vascular plants endemic to Yakushima Island, Japan. *Journal of Faculty of Science University of Tokyo III*, 14: 69-111.