

VEGETATIVE PROPAGATION OF DIFFERENT CYCLAMEN SPECIES VIA ADVENTITIOUS SHOOT FORMATION FROM SEEDLING TISSUE

Anika Nadja Sabine Prange^{1*}, Margrethe Serek¹, and Traud Winkelmann²

¹Gottfried Wilhelm Leibniz University of Hannover, Institute of Floriculture and Woody Plant Science, Floriculture Section, 2 Herrenhaeuser str., D-30419 Hannover, Germany,

*Fax: 0049 511 762 2654, *E-mail: prange@zier.uni-hannover.de

²University of Applied Sciences Weihenstephan and Research Station for Horticulture, Am Staudengarten 8, D-85354 Freising, Germany

REFERENCES

- Agresti A. (1990). Categorical data analysis. John Wiley & Sons, Inc., New York: 59-64.
- Bellamine J., Penel C., Greppin H., Gaspar T. (1998). Confirmation of the role of auxin and calcium in the late phases of adventitious root formation. *Plant Growth Regulation*, 26: 191-194.
- Ewald A. (1996). Interspecific hybridisation between *Cyclamen persicum* Mill. and *C. purpurascens* Mill. *Plant Breeding*, 115: 162-166.
- Geier T., Kohlenbach H. W., Reuther G. (1990). Cyclamen – history and economic importance. In: Ammirato P. V., Evans D. R., Sharp W. R., Bajaj Y. P. S. (Eds.). *Handbook of Plant Cell Culture, Vol. 5 Ornamental Species*. McGraw-Hill Publishing Company, New York: 352-374.
- Glasau F. (1939). Monographie der Gattung *Cyclamen* auf morphologisch-cytologischer Grundlage. *Planta*, 30: 507-550 (in German).
- Goebel K. (1908). Einleitung in die experimentelle Morphologie der Pflanzen. B. G. Teubner, Leipzig, 260 pp. (in German).
- Hawkes H. Y., Wainwright H. (1987). *In vitro* organogenesis of *Cyclamen persicum* Mill. seedling tissue. *Acta Horticulturae*, 212: 711-714.
- Hildebrand F. (1898). Die Gattung *Cyclamen*, eine Systematische und Biologische Monographie. Fischer, Jena, 190 pp. (in German).
- Hill A. W. (1920). Studies in seed germination: Experiments with *Cyclamen*. *Annals of Botany*, 34: 417-429.
- Ishizaka H. (1996). Interspecific hybrids of *Cyclamen persicum* and *C. graecum*. *Euphytica*, 91: 109-17.
- Ishizaka H., Uematsu J. (1992). Production of interspecific hybrids of *Cyclamen persicum* Mill. and *C. hederifolium* Aiton. by ovule culture. *Japanese Journal of Breeding*, 42: 353-366.
- Ishizaka H., Uematsu J. (1995). Interspecific hybrids of *Cyclamen persicum* Mill. and *C. purpurascens* Mill. produced by ovule culture. *Euphytica*, 82: 31-37.
- Karam N. S., Al-Majathoub M. (2000a). *In vitro* shoot regeneration from mature tissue of wild *Cyclamen persicum* Mill. *Scientia Horticulturae*, 86: 323-333.
- Karam N. S., Al-Majathoub M. (2000b). Direct shoot regeneration and microtuberisation in wild *Cyclamen persicum* Mill using seedling tissue. *Scientia Horticulturae*, 86: 325-346.
- Murashige T., Skoog F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- Schwenkel H. G., Winkelmann T. (1998). Plant regeneration via somatic embryogenesis from ovules of *Cyclamen persicum* Mill. *Plant Tissue Culture and Biotechnology*, 4: 28-34.
- Sharma D. R., Kaur R., Kumar K. (1996). Embryo rescue in plants – a review. *Euphytica*, 89: 325-337.
- Winkelmann T., Specht J., Serek M. (2006). Efficient plant regeneration from protoplasts isolated from embryogenic suspension cultures of *Cyclamen persicum* Mill. *Plant Cell, Tissue and Organ Culture*, 86: 337-347.
- Winkler H. (1902) Über die Regeneration der Blattspreite bei einigen *Cyclamen* Arten. *Berichte der Deutschen Botanischen Gesellschaft*, 20: 81-87 (in German).