

MICROPROPAGATION OF *PHALAENOPSIS*-HYBRIDS IN TEMPORARY IMMERSION SYSTEM (TIS) – EFFECTS OF EXUDATED PHENOLIC SUBSTANCES ON PLANT DEVELOPMENT

Claudia Pisowotzki*, Katja Saare-Surminski, and Reinhard Lieberei

Department of Crop Plants and Plant Ecology, Biocentre Klein Flottbek and Botanical Garden,
University of Hamburg, 18 Ohnhorst str., 22609 Hamburg, Germany,
*Fax: + 49 40 42816 565, *E-mail: claudiapisowotzki@alice-dsl.de

REFERENCES

- Bhatt I. D., Dhar U. (2004). Factors controlling micropropagation of *Myrica esculenta* Buch.- Ham. ex D. Don: A high value wild edible of Kumaun Himalaya. *African Journal of Biotechnology*, 3(10): 534-540.
- Blum U., Gerig T. M. (2005). Relationships between phenolic acid concentrations, transpiration, water utilization, leaf area expansion, and uptake of phenolic acids: Nutrient culture studies. *Journal of Chemical Ecology*, 31 (8): 1907-1932.
- Chen Y.-C., Chang C., Chang W.-C. (2000). A reliable protocol for plant regeneration from callus culture of *Phalaenopsis*. *In Vitro Cellular and Developmental Biology Plant*, 36: 420-423.
- Frazier R. A., Papadopoulou A., Mueller-Harvey I., Kisson D., Green R. J. (2003). Probing Protein-Tannin Interactions by Isothermal Titration Microcalorimetry. *Journal of Agricultural and Food Chemistry*, 51: 5189-5195.
- Hempfling T., Preil W. (2005). Application of a temporary immersion system. *In: Hroslef-Eide A. K., Preil W. (Eds.). Liquid culture systems for in vitro plant propagation.* Springer, Berlin: 197-211.
- Lorenzo J. C., Angeles Blanco M. de los, Pelaez O., Gonz alez A., Cid M., Iglesias A., Gonz lez B.; Escalona M., Espinosa P., Borroto C. (2001). Sugarcane micropropagation and phenolic excretion. *Plant Cell, Tissue and Organ Culture*, 65: 1-8.
- Murashige T., Skoog F. (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.