

ROOTING, QUANTIFICATION OF INDOLE-3-ACETIC ACID AND INDOLE-3-BUTYRIC ACID, AND IBA TRANSPORT IN *GREVILLEA*

Santi Krisantini^{1*}, Margaret Johnston², Christine Anne Beveridge³, John Ross⁴, and Richard Williams²

¹Department of Agronomy and Horticulture, Bogor Agricultural University, Bogor, Indonesia 16680,

*Fax: + 62-251-629 353, *E-mail: krisantini@yahoo.com.au

²School of Land, Crop and Food Sciences, University of Queensland, Gatton Campus,
Queensland, Australia 4343

³School of Integrative Biology, University of Queensland, St. Lucia Campus, Queensland, Australia 4071

⁴School of Plant Science, The University of Tasmania, Australia 7001

REFERENCES

- BARALDI R., BERTAZZA G., PREDIERI S., BREGOLI A. M., COHEN J. D. (1993). Uptake and metabolism of indole-3-butyric acid during the *in-vitro* rooting phase in pear cultivars (*Pyrus communis*). *Acta Horticulturae*, 329: 289-291.
- BAUER L. M. (1999). The Propagation, growth and development of *Persoonia virgata*. PhD Thesis. The University of Queensland, Gatton, 207 pp.
- BEVERIDGE C. A., ROSS J. J., MURFET I. C. (1994). Branching mutant *rms-2* in *Pisum sativum*: grafting studies and endogenous indole-3-acetic acid levels. *Plant Physiology*, 104: 953-959.
- BEVERIDGE C. A., SYMONS G. M., ROSS J. R., RAMEAU C. (1997). The *rms 1* mutant of pea has elevated indole-3-acetic acid levels and reduced root-sap zeatin riboside content but increased branching controlled by graft-transmissible signals. *Plant Physiology*, 116: 1251-1258.
- BOERJAN W., CERVERA M. T., DELARUE M., BEECKMAN T., DEWITTE W., BELLINI C., CABOCHE M., ONCKELEN H. V., VAN MONTAGU M., INZE D. (1995). Superroot, a recessive mutation in *Arabidopsis*, confers auxin overproduction. *The Plant Cell*, 7: 1405-1419.
- BROWN D. E., RASHOTTE A. M., MURPHY A. S., NORMANLY J., TAGUE B. W., PEER W. A., TAIZ L., MUDAY G. K. (2001). Flavonoids act as negative regulators of auxin transport *in vivo* in *Arabidopsis*. *Plant Physiology*, 126: 524-535.
- BURGESS J. (1985). Polarity and development. *In: An introduction to plant cell development*. Cambridge University Press, UK: 203-206.
- CABONI E., LAURI P., TONELLI M. G., LACOVACCI P., DAMIANO C. (1997). Biochemical and molecular factors affecting *in vitro* rooting ability in almond. *In: Altman A., Waisel W. (Eds). Biology of Root Formation and Development*, Plenum Press, New York: 117-124.
- CHHUN T., TAKETA S., ICHII M., TSURUMI S. (2004). Different behaviour of indole-3-acetic acid and indole-3-butyric acid in stimulating lateral root development in rice (*Oryza sativa* L.). *Plant Growth Regulation*, 43: 135-143.
- CHHUN T., TAKETA S., ICHII M., TSURUMI S. (2005). Involvement of ARM2 in the uptake of indole-3-butyric acid in rice (*Oryza sativa* L.) roots. *Plant Cell Physiology*, 46: 1161-1164.
- CHHUN T., UNO Y., TAKETA S., AZUMA T., ICHII M., OKAMOTO T., TSURUMI S. (2007). Saturated humidity accelerates lateral root development in rice (*Oryza sativa* L.) seedlings by increasing phloem-based auxin transport. *Journal of Experimental Botany*, 58: 1695-1709.
- COLLET D. (1991). Models for Binary and Binomial Data. *In: Collett D. (Ed.). Modelling Binary Data*, Chapman and Hall/CRC: 43-92.
- DAVIES P. J. (1995). The plant hormones: their nature, occurrence and functions. *In: Davies P. J. (Ed.). Plant Hormone*, Kluwer Academic Press, London, UK: 1-12.
- DE BILLY F., GROSJEAN C., MAY S., BENNETT M., CULLIMORE J. V. (2001). Expression studies on AUX1-like genes in *Medicago truncatula* suggest that auxin is required at two steps in early nodule development. *Molecular Plant-Microbe Interactions*, 14: 267-277.
- DE KLERK G.-J., VAN DER KRIEKEN W., DE JONG J. C. (1999). Review: the formation of adventitious roots, new concepts, new possibilities. *In Vitro Cellular and Developmental Biology-Plant*, 35: 189-199.
- DIAZ-SALA C., HUTCHISON K. W., GOLDFARB B., GREENWOOD M. S. (1998). Maturation-related loss in rooting competence by loblolly pine stem cuttings: the role of auxin transport, metabolism and tissue sensitivity. *Physiologia Plantarum*, 97: 481-490.
- EPSTEIN E., ACKERMAN A. (1993). Transport and metabolism of indole-3-butyric acid in cuttings of *Leucadendron discolor*. *Plant Growth Regulation*, 12: 17-22.
- EPSTEIN E., LUDWIG-MULLER J. (1993). Indole-3-butyric acid in plants: occurrence, synthesis, metabolism and transport. *Physiologia Plantarum*, 88: 382-389.

- FORD Y. Y., BONHAM E. C., CAMERON R. W. F., BLAKE P. S., JUDD H. L., HARRISON-MURRAY R. S. (2002). Adventitious rooting: examining the role of auxin in an easy- and difficult-to-root plant. *Plant Growth Regulation*, 36: 149-159.
- JONES S. E., DEMEO J. S., DAVIES N. W., NOONAN S., ROSS J. J. (2005). Stems of the *Arabidopsis* pin1-1 mutant are not deficient in free indole-3-acetic acid. *Planta*, 222: 530-534.
- KIBBLER H. (2001). Barriers in adventitious root formation in *Baccharis citriodora*. PhD Thesis. School of Agriculture and Horticulture. The University of Queensland, Gatton, 201 pp.
- KRISANTINI S., JOHNSTON M., WILLIAMS R. R., BEVERIDGE C. A. (2003). Propagation of *Grevillea*. Combined Proceedings of International Plant Propagators' Society, 53: 154-158.
- LANTERI M. L., PAGNUSSAT G. C., LAMATTINA L. (2006). Calcium and calcium-dependent protein kinases are involved in nitric oxide- and auxin-induced adventitious root formation in cucumber. *Journal of Experimental Botany*, 57: 1341-1351.
- Liu J., Mukherjee I., Reid D. M. (1990). Adventitious rooting in hypocotyls of sunflower (*Helianthus annuus*). *Physiologia Plantarum*, 78: 268-276.
- LUDWIG-MULLER J. (2003). Peroxidase isoenzymes as markers for the rooting ability of easy-to-root and difficult-to-root *Grevillea* species and cultivars of *Protea obtusifolia* (Proteaceae). *In Vitro Cellular and Developmental Biology-Plant*, 39: 377-383.
- LUDWIG-MULLER J., COHEN D. (2002). Identification and quantification of three active auxins in different tissues of *Tropaeolum majus*. *Physiologia Plantarum*, 115: 320-329.
- LUDWIG-MULLER J., EPSTEIN E. (1993). Analysis of indole-3-butyric acid in *Arabidopsis thaliana*. *Acta Horticulturae*, 329: 109-111.
- LUDWIG-MULLER J., VERTOCNIK A., TOWN C. D. (2005). Analysis of IBA-induced adventitious root formation on *Arabidopsis* stem segments. *Journal of Experimental Botany*, 56: 2095-2105.
- MARKS T. R., FORD Y. Y., CAMERON R. W. F., GOODWIN C., MYERS P. E., JUDD H. L. (2002). A role for polar auxin transport in rhizogenesis. *Plant Cell, Tissue and Organ Culture*, 70: 189-198.
- NAG S., SAHA K., CHOUDHURI M. A. (2001). Role of auxin and polyamines in adventitious root formation in relation to changes in compounds involved in rooting. *Journal of Plant Growth Regulation*, 20: 182-194.
- NORDSTROM A. C., JACOBS F. A., ELIASSON L. (1991). Effect of exogenous indole-3-acetic acid and indole-3-butyric acid on internal levels of the respective auxins and their conjugation with aspartic acid during adventitious root formation in pea cuttings. *Plant Physiology*, 96: 856-861.
- RASHOTTE A. M., POUPART J., WADDELL C. S., MUDAY G. K. (2003). Transport of the two natural auxins, indole-3-butyric acid and indole-3-acetic acid, in *Arabidopsis*. *Plant Physiology*, 133: 761-772.
- SCHWARZ J. L., GLOCKE P. L., SEDGLEY M. (1999). Adventitious root formation in *Acacia baileyana* F. Muell. *Journal of Horticultural Sciences and Biotechnology*, 74: 561-565.
- SINGH D. (2000). *Grevillea* Hybrids: 'Coastal Dawn'. *Plant Varieties Journal*, 13: 25-26
- SUTTER E. G., COHEN J. D. (1992). Measurement of indolebutyric acid in plant tissues by isotope dilution gas chromatography-mass spectrometry analysis. *Plant Physiology*, 99: 1719-1722.
- SYMONS G. M., ROSS J. J., MURFET I. C. (2002). The *bushy* pea mutant is IAA-deficient. *Physiologia Plantarum*, 116: 389-397.