Propagation of Ornamental Plants
Vol. 9, № 4, 2009: 207-215

USE OF THE PMI/MANNOSE SELECTION SYSTEM FOR GENETIC TRANSFORMATION OF *KALANCHOE BLOSSFELDIANA* POELLN.

Agnieszka Ilczuk[1,2], Heiko Mibus[1], Malthe Schmidt[1], and Margrethe Serek[1]*

[1]Faculty of Natural Sciences, Institute for Ornamental and Woody Plant Science,
Leibniz University of Hannover, 2 Herrenhaeuser str., 30419 Hannover, Germany,
*Fax: + 49 511 762 26 54, *E-mail: serek@zier.uni-hannover.de
[2]Department of Ornamental Plants, Faculty of Horticulture and Landscape Architecture,
Warsaw University of Life Sciences, 166 Nowoursinowska str., 02-787 Warsaw, Poland

REFERENCES

Aida R., Shibata M. (1996). Transformation of *Kalanchoe blossfeldiana* mediated by *Agrobacterium tumefaciens* and transgene silencing. Plant Science, 121:175-185.

Bailey M. A., Kaeppler H. F. (2001). Special-workshop: alternative markers for plant transformation. Workshop presentations from the 2000 World Congress on *In Vitro* Biology. In vitro Cellular & Development Biology-Plant, 37: 101-102.

Boscariol R. L., Almeida W. A. B., Derbyshire M. T. V. C., Mourao Filho F. A. A., Mendes B. M. J. (2003). The use of the PMI/mannose selection system to recover transgenic sweet orange plants (*Citrus sinensis* L. Osbeck). Plant Cell Reports, 22: 122-128.

Brandt P. (2001) Die neue Generation von Marker-Genen zum frühen Erkennen von genetisch veränderten Pflanzen. Bundesgesundheitsbl-Gesundheitsforsch-Gesundheitsschutz, 44: 1188-1193.

Dale E. C., Ow D. W. (1991). Gene transfer with subsequent removal of the selection gene from the host genome. Proceedings of the National Academy of Sciences of USA, 88: 10558-10562.

De la Riva G. A., Cabrera J. G., Padron R. V., Padro C. A. (1998). *Agrobacterium tumefaciens*: a natural tool for plant transformation. European Journal of Biotechnology, 1: 1-18.

Degenhardt J., Poppe A., Montag J., Szankowski I. (2006). The use of phosphomannose-isomerase/mannose selection system to recover transgenic apple plants. Plant Cell Reports, 25: 1149-1156.

Ding Z. S., Zhao M., Jing Y. X., Li L. B., Kuang T. Y. (2006). Efficient *Agrobacterium*-mediated transformation of rice by phosphomannose isomerase/mannose selection. Plant Molecular Biology Reporter, 24: 295-303.

Flavell R. B., Dart E., Fuchs R. L., Fraley R. T. (1992). Selectable marker genes: safe for plants? Biotechnology, 10: 141-144.

Gamborg O. L., Miller R. A., Ojima K. (1968). Nutrient requirements of suspension culture of soybean root cells. Experimental Cell Research, 50: 151-158.

Ghorbel R., La-Malfa S., Lopez M. M., Petit A., Navarro L., Pena L. (2001). Additional copies of *virG* form pTiBo542 provide a super-transformation ability to *Agrobacterium tumefaciens* in citrus. Physiological and Molecular Plant Pathology, 58: 103-110.

Goldsworthy A., Street H. E. (1965). The carbohydrate nutrition of tomato roots VIII. The mechanism of the inhibition by D-mannose of the respiration of excised roots. Annals of Botany, 29: 45-58.

Haldrup A., Petersen S. G., Okkels F. T. (1998a). The xylose isomerase gene from *Thermoanaerobacterium thermosulfurogenes* allows effective selection of transgenic plants cells using D-xylose as the selection agent. Plant Molecular Biology, 37: 287-296.

Haldrup A., Petersen S. G., Okkels F. T. (1998b). Positive selection: A plant selection principle based on xylose isomerase, an enzyme used to the food industry. Plant Cell Reports, 18: 76-81.

He Z. Q., Duan Z., Liang W., Chen F., Yao W., Liang H., Yue Ch., Sun Z., Chen F., Dai J. (2006). Mannose selection system used for cucumber transformation. Plant Cell Reports, 25: 953-958.

He Z. Q., Fu Y. P., Si H. M., Hu G. Ch., Zhang S. H., Yu Y. H., Sun Z. X. (2004). Phosphomannose isomerase (*pmi*) gene as a selectable marker for rice transformation via *Agrobacterium*. Plant Science, 166: 17-22.

Hoa T. T. C., Bong B. B. (2002). *Agrobacterium*-mediated transformation of rice embryogenic suspension cells using phosphomannose isomerase gene, *pmi*, as a selectable marker. Omonrice, 10: 1-5.

Hood E. E., Gelvin S. B., Melchers L. S., Hoekema A. (1993). New *Agrobacterium* helper plasmids for gene transfer to plants. Transgenic Research, 2: 208-212.

Jain M., Chenegalrayan K., Abouzid A., Gallo M. (2007). Prospecting the utility of a PMI/mannose selection system for the recovery transgenic sugarcane (*Saccharum* spp. hybrid) plants. Plant Cell Reports, 26: 581-590.

Janakiraman V., Steinau M., McCoy S., Trick H. N. (2002). Recent advances in wheat transformation. In Vitro Cellular & Developmental Biology-Plant, 38: 404-414.

Jefferson R. A., Kavanagh T. A., Bevan M. W. (1987). GUS fusions: ß-glucoronidase as a sensitive and versatile gene fusion marker in higher plants. EMBO Journal, 6: 3901-3907.

Joersbo M. (2001). Advances in the selection of transgenic plants using non-antibiotic marker genes. Physiologia Plantarum, 111: 269-272.

Joersbo M., Donaldson I., Kreiberg J., Petersen S.G., Brunstedt J., Okkels F. T. (1998). Analysis of mannose selection used for transformation of sugar beet. Molecular Breeding, 4 (2): 111-117.

Joersbo M., Okkels F. T. (1996). A novel principle for selection of transgenic plant cells: positive selection. Plant Cell Reports, 16 (3/4): 219-221.

Kim J. Y., Jung M., Kim H. S., Lee Y. H., Choi S. H., Lim Y. P., Min B. W., Yang S. G., Harn C. H. (2002). A new selection system for pepper regeneration by mannose. Journal of Plant Biotechnology, 4: 129-134.

Lamblin F., Aime A., Hano C., Roussy I., Domon J. M., Van-Droogenbroeck B., Laine E. (2007). The use of the phosphomannose isomerase gene as alternative selectable marker for *Agrobacterium*-mediated transformation of flax (*Linum usitatissimum*). Plant Cell Reports, 26: 765-772.

Li H. Q., Kang P. J., Li M. L., Li M. R. (2007). Genetic transformation of *Torenia fournieri* using the PMI/mannose selection system. Plant Cell, Tissue and Organ Culture, 90: 103-109.

Lin H. Y., Garcia-Assad N., Kuo J. (1994). Effect of *Agrobacterium* cell concentration on the transformation efficiency of tabacco and *Arabidopsis thaliana*. Focus, 16: 72-77.

Murashige T., Skoog F. (1962). A revised medium for rapid growth and bioassays with tabacco tissue cultures. Physiologia Plantarum, 15: 473-497.

Nap J. P., Bijvoet J., Stiekema W. J. (1992). Biosafety of kanamycin resistant transgenic plants. Transgenic Research, 1: 239-249.

Negretto D., Jolley M., Beer S., Wenck A. R., Hansen G. (2000). The use of phosphomannose isomerase as a selectable marker to recover transgenic maize plants (*Zea mays* L.) via Agrobacterium transformation. Plant Cell Reports, 19: 798-803.

Okkels F. T., Ward J. L., Joersbo M. (1997). Synthesis of cytokinin glucuronides for the selection of transgenic plant cells. Phytochemistry, 46: 801-804.

Orlikowska T. (2001). Influence of growrth regulators and antibiotics on regeneration of *Kalanchoe blossfeldiana* Poelln. *in vitro*. Biology Bulletin of Poznan, 38: 29-34.

Peng S. Q., Chen S. C. (2005). A novel selection system for tomato transformation by mannose. Journal of Agricultural Biotechnology, 13: 141-144.

Penna S., Sagi L., Swennen R. (2002). Positive selectable marker genes for routine plant transformation. In Vitro Cellular & Developmental Biology-Plant, 38: 125-128.

Ramesh S. A., Kaiser B. N., Franks T., Collins G., Sedgley M. (2006). Improved methods in *Agrobacterium*-mediated transformation of almond using positive (mannose/*pmi*) or negative (kanamycin resistance) selection-based protocols. Plant Cell Reports, 25: 821-828.

Reed J., Privalle L., Powell M. L., Meghji M., Dawson J., Dunder E., Suttie J., Wenck A., Launis K., Kramer C., Chang Y. F., Hansen G., Wright M. (2001). Phosphomannose isomerase: an efficient selectable marker for plant transformation. In Vitro Cellular & Developmental Biology-Plant, 37: 127-132.

Rieger M., Preston C., Powles S. (1999). Risks of gene flow from transgenic herbicyde-resistant canola (*Brassica napus*) to weedy relatives in southern Australian crooping systems. Australian Journal of Agricultural Research, 50: 115-128.

Saini R., Jaiwal P. K. (2007). *Agrobacterium tumefaciens*-mediated transformation of blackgram: an assessment of factors influencing the efficiency of *uid*A gene transfer. Biologia Plantarum, 51: 69-74.

Sanikhani M., Mibus H., Stummann B. M., Serek M. (2007). *Kalanchoe blossfeldiana* plants expressing the *Arabidopsis etr1-1* allele show reduced ethylene sensitivity. Plant Cell Reports, 27: 729-737.

Seitz C., Bruna S., Li H., Hauser B., Forkmann G. (2007). Establishment of a mannose-based selection system for the transformation of *Torenia* hybrids. Acta Horticulturae, 743: 139-142.

Sheikholeslam S. N., Weeks D. P. (1987). Acetosyringone promoter high efficiency transformation of *Arabidopsis thaliana* explants by *Agrobacterium tumefaciens*. Plant Molecular Biology, 8: 291-298.

Shi-Rong J., Mei-Zhu Y., Russ O., Nam-Hai Ch. (1989). High frequency transformation of *Kalanchoe laciniata*. Plant Cell Reports, 8: 336-340.

Sriskandarajah S., Frello S., Jorgensen K., Serek M. (2004). *Agrobacterium tumefaciens*-mediated transformation of *Campanula carpatica*: factors affecting transformation and regeneration of transgenic shoots. Plant Cell Reports, 23: 59-63.

Stachel S.E., Nester E. W., Zambryski P. (1986). A plant cell factor induces *Agrobacterium tumefaciens* vir gene expression. Proceedings of the National Academy of Sciences of USA, 83: 379-383.

Stevis P. E., Ho N. W. Y. (1987). Positive selection vectors based on xylose utilization suppression. Gene, 55: 67-74.

Stevis P. E., Ho N. W. Y. (1988). Short communication. A novel *xyl*B-based positive selection vector. Plasmid, 20: 92-95.

Teixeira da Silva J. A.(2004). Evaluation of carbon sources as positive selection agents for chrysanthemum (*Dendranthema* x *grandiflorum*) transformation. New Zealand Journal of Crop and Horticultural Science, 32: 55-67.

Todd R., Tague B. W. (2001). Phosphomannose isomerase: a versatile selectable marker for *Arabidopsis thaliana* germ-line transformation. Plant Molecular Biology Reporter, 19: 307-319.

Zai-Song D., Ming Z., Yu-Xiang J., Liang-Bi L., Ting-Yun K. (2006). Efficient *Agrobacterium*-mediated transformation of rice by phosphomannose isomerase/mannose selection. Plant Molecular Biology Reporter, 24: 295-303.

Zhu L. H., Li X. Y., Ahlman A., Xue Z. T., Welander M. (2004). The use of mannose as a selection agent in transformation of the apple rootstock M26 via *Agrobacterium tumefaciens*. Acta Horticulturae, 663: 503-506.